

Some aspects of a vector basis for a controllable group system, group code, and group shift

K. M. Mackenthun Jr. (email: ken1212576@gmail.com)

November 13, 2012

ABSTRACT

In a group trellis, the sequence of branches that split from the identity path and merge to the identity path form two normal chains. The Schreier refinement theorem can be applied to these two normal chains to obtain a Schreier series, a normal chain of the branch group B of the group trellis. It is shown that the components of the shortest length generator sequences of Forney and Trott form a complete system of coset representatives for the Schreier series decomposition of B . These components can be used to construct a natural time domain encoder with the form of a time convolution. The Schreier series and encoder have a natural shift structure related to B , and we find its graph automorphism group. Using the natural shift structure, we find a vector basis group for B which is isomorphic to B . When the basis is formed using generator sequences, the graph automorphism group gives a bound on the number of generator bases. Using the graph automorphism group, the encoder here is compared to an encoder of Forney and Trott. When B is abelian, there is a symmetry and duality between the two encoders which has a transform domain interpretation between a time and spectral domain. When B is nonabelian, the comparison shows that the components of the generator sequences must obey commutative restrictions, and so it is inherently difficult for a branch group to be nonabelian.

1. INTRODUCTION

The idea of group shifts and group codes is important in several areas of mathematics and engineering such as symbolic dynamics, linear systems theory, and coding theory. Research in this area started with the work of Kitchens [1], Willems [2], Forney and Trott [3], and Loeliger and Mittelholzer [4].

Kitchens [1] introduced the idea of a group shift [12] and showed that a group shift has finite memory, i.e., it is a shift of finite type [12]. A group shift is a fundamental example of a time invariant group code [3], also called a time invariant group system [3]. Forney and Trott [3] showed that any time invariant group code is equivalent to a labeled group trellis section. They showed that any group code that is complete (any global constraints can be determined locally, see [3]) is equivalent to a sequence of connected labeled group trellis sections (which may vary

in time). They explained the important idea of shortest length code sequences, or generator sequences. At each time epoch, a finite set of generator sequences can be used to construct a “local” section of the code. Using the generator sequences, any group code can be mechanized with a minimal encoder which has a shift register structure.

Forney and Trott start with the most general setting, a set of sequences with a group property, a group code, and then analyze further to determine the state structure, group trellis section, group trellis, encoder structure, and other properties of the code; the related work of [2] in systems theory also uses this same top down approach.

The work of Loeliger and Mittelholzer [4] starts with a group trellis. Among other results, Loeliger and Mittelholzer give an abstract characterization of groups which can be the branch group B of a group trellis. They also give a shift register structure for a group trellis, which uses the Forney-Trott approach modified to a group trellis. Instead of using groups related to code sequences as in [3], their encoder is described using groups related to their abstract characterization of a branch group B . Aside from this abstract description, the encoder of [4] is essentially the Forney-Trott encoder applied to a group code which consists of code sequences of branches from a group trellis. Thus we call the encoder in [4] the *branch group version of the Forney-Trott encoder*.

This paper also gives an abstract characterization of B , but it is different from [4]. In a group trellis, the sequence of branches that split from the identity path and merge to the identity path form two normal chains, $\{X_j\}$ and $\{Y_i\}$, respectively. In this paper we apply the Schreier refinement theorem to $\{X_j\}$ and $\{Y_i\}$. The refinement of the two normal chains gives another normal chain, the Schreier series, which is also a normal chain of B . The Schreier series can be written in the form of a matrix, with rows and columns determined by $\{X_j\}$ and $\{Y_i\}$. When the group trellis is controllable, the matrix reduces to a triangular form, called the controllable matrix. The controllable matrix is an echo of matrix ideas used in classical linear systems analysis.

In this paper, we start with a group trellis as in [4]. However we feel the approach here also suffices when starting with a group code as in [3]. The only change required is to replace the Schreier series obtained from the Schreier refinement of trellis normal chains $\{X_j\}$ and

$\{Y_i\}$ with an analogous normal chain which formally replaces $\{X_j\}$ and $\{Y_i\}$ with corresponding sequences in the group code.

Using properties of the controllable matrix and Schreier series, we show that components of the shortest length generator sequences of Forney and Trott form a complete system of coset representatives for the Schreier series decomposition of the branch group B of the code given by the controllable matrix. This shows the generator sequences are intimately related to the branch group B .

Based on the controllable matrix, we give a minimal encoder structure for a group code which is an estimator. As in [3, 4], the encoder uses shortest length generator sequences, but here the components of the generator sequences give a time domain convolution. Therefore this encoder appears to be a natural time domain form for a group code encoder. The encoder has a register implementation which is somewhat different from the classical shift register structure.

The Schreier series and encoder have a natural shift structure related to B , and we find its graph automorphism group. Using the natural shift structure, we find a vector basis group for B which is isomorphic to B . When the basis is formed using generator sequences, the graph automorphism group gives a bound on the number of generator bases. The encoder here is compared to an encoder of Forney and Trott. When B is abelian, there is a symmetry and duality between the two encoders which has a transform domain interpretation between a time and spectral domain. When B is nonabelian, the comparison shows that the components of the generator sequences must obey commutative restrictions, and so it is inherently difficult for a branch group to be nonabelian.

The recent work of Forney [7] discusses a basis for a group code or group system which uses generator sequences. Here we give a vector basis whose components are not sequences but individual branches. Although components of the generator sequences can be used to form a vector basis, it is not necessary to use generators to form a vector basis.

Forney and Trott [3] have shown that any group system can be reduced to a group trellis of concatenated trellis sections, composed of states and branches. The states have the properties of states in linear systems theory [2]. This work gives a vector representation of the states and branches. The vector representation is essentially derived from the behavior of the trellis. Thus the states and branches are essentially intertwined with the trellis. Therefore the notions of trellis, branches, and states are not fundamentally separate, but they are just different ways of expressing the same underlying behavioral system. For example, two different states are not really distinct but share some degree of common behavior which can be made precise by examining their basis vectors.

This paper is organized as follows. Section 2 defines

a group trellis section and group trellis. We study an ℓ -controllable group trellis, in which each state can be reached from any other state in ℓ branches. Section 3 defines the Schreier series. For an ℓ -controllable group trellis, the Schreier series can be arranged in the form of a triangular matrix, called the controllable matrix. In Section 4, we analyze the structure of the controllable matrix. We show that the rows of the controllable matrix form quotient groups, and the generator sequences of Forney and Trott are a transversal of the quotient groups. This means the components of the generators form a complete system of coset representatives for the Schreier series decomposition of B . The same system of coset representatives can be used for the Schreier series decomposition of the time reversed group trellis. Based on the analysis in Section 4, Section 5 gives an encoder for the group trellis. The encoder uses shortest length generator sequences and has a time convolution property.

The controllable matrix and encoder have a natural shift structure. In Section 6 we identify this shift structure, and find its graph automorphism group. We find a set of separating permutations which completely specifies the automorphism group. This allows us to count the number of automorphisms. In Section 7, we discuss a vector basis formed using factor groups of the finite chain $\{X_j\}$. The vector basis uses representatives of cosets in the factor groups, and each choice of representatives gives a different basis. We show that B is isomorphic to a group of equivalence classes of vector expansions of g in each basis, for each $g \in B$. A generator basis can be formed using components of generators. We give a change of basis theorem for the generator bases. Using the graph automorphism group, we bound the number of generator bases. Finally we compare our encoder with the branch group version of the Forney-Trott encoder. The two encoders only differ by an interchange of inner and outer product, or a rearrangement of group elements in a product. Since the two encoders must have isomorphic graphs, and since the same generators can be used in both encoders, there are commutative restrictions on components of the generators. Therefore any nonabelian group must satisfy certain commutative restrictions. When B is abelian, the two encoders have a symmetry and duality, which can be interpreted as a transform relation between a time and spectral domain.

2. GROUP CODE AND GROUP TRELLIS

As in [11], given groups G_1, G_2, G_3 , we say G is a *subdirect product* of G_1, G_2 , and G_3 if it is a subgroup of $G_1 \times G_2 \times G_3$ and each coordinate of G takes all values in its respective group G_1, G_2 , or G_3 . We also say G is a subdirect product of $G_1 \times G_2 \times G_3$.

Definition. A *group trellis section* is a subdirect product B of $\Sigma \times A \times \Sigma$. We call B the *branch group* and Σ the *state group*. The left states of B, B^- , are the group Σ , the first coordinate of $\Sigma \times A \times \Sigma$. The right states of B ,

B^+ , are the group Σ , the last coordinate of $\Sigma \times A \times \Sigma$. We call A the *label group* or *alphabet*. The elements of B are *branches* b , where $b = (s, a, s') \in \Sigma \times A \times \Sigma$. The left state of b , b^- , is s , and the right state of b , b^+ , is s' . •

We can think of a group trellis section B as a bipartite graph with branches in B and vertices in Σ , where there is a branch (s, a, s') with label a between two vertices s and s' if and only if $(s, a, s') \in B$.

Consider the projection map $\pi^- : B \rightarrow B^- = \Sigma$ given by the assignment $(s, a, s') \mapsto s$. This is a homomorphism with kernel X_0 , the subgroup of all elements of B of the form $(\mathbf{1}_\Sigma, a, s')$, where $\mathbf{1}_\Sigma$ is the identity of Σ . Then the branches that split from each left state of B^- are a coset of X_0 , and $B/X_0 \simeq B^-$. Consider the projection map $\pi^+ : B \rightarrow B^+ = \Sigma$ given by the assignment $(s, a, s') \mapsto s'$. This is a homomorphism with kernel Y_0 , the subgroup of all elements of B of the form $(s, a, \mathbf{1}_\Sigma)$. Then the branches that merge to each right state of B^+ are a coset of Y_0 , and $B/Y_0 \simeq B^+$.

In general, for a subset or subgroup of branches $G \subset B$, define G^- to be the left states of G , and define G^+ to be the right states of G .

Proposition 1 *Let G be a subgroup of B . We have $G \cap X_0 \triangleleft G$ and $G \cap Y_0 \triangleleft G$. The branches that split from each left state of G , G^- , are a coset of $G \cap X_0$, and $G^- \simeq G/G \cap X_0$. The branches that merge to each right state of G , G^+ , are a coset of $G \cap Y_0$, and $G^+ \simeq G/G \cap Y_0$.*

We think of a *group trellis* as a connected sequence of group trellis sections B [3]. The branches of the group trellis occur at time epochs t , which are integers in the range $-\infty < t < \infty$. Let C be a group trellis, and let $\mathbf{c}$ be a trellis path in C . Then

$$\mathbf{c} = \dots, b_{-1}, b_0, b_1, \dots, b_t, \dots,$$

where component b_t is a *branch at time epoch t* , and $b_t \in B$. Define the projection map at time t , $\chi_t : C \rightarrow B$, by the assignment $\mathbf{c} \mapsto b_t$. Define the projection map $\chi_{[t_1, t_2]} : C \rightarrow B \times \dots \times B$ by the assignment $\mathbf{c} \mapsto (b_{t_1}, \dots, b_{t_2})$, the components of $\mathbf{c}$ over the time interval $[t_1, t_2]$. We say that $(b_{t_1}, \dots, b_{t_2})$ is a *trellis path segment of length $t_2 - t_1 + 1$, on time interval $[t_1, t_2]$* .

Define C^+ to be the set of all codewords for which $b_t = \mathbf{1}$ for $t < 0$, where $\mathbf{1}$ is the identity of B . Define C^- to be the set of all codewords for which $b_t = \mathbf{1}$ for $t \geq 0$. Define $C_{[t_1, t_2]}$ to be the set of all codewords which are the identity outside the time interval $[t_1, t_2]$; this is a group. We say that a codeword $\mathbf{c} \in C_{[t_1, t_2]}$ has *span $t_2 - t_1 + 1$* if $c_{t_1} \neq \mathbf{1}$ and $c_{t_2} \neq \mathbf{1}$. Similary define $C_{[t_1, t_2]}$ and $C_{(t_1, t_2)}$.

For all integers j , define

$$X_j \stackrel{\text{def}}{=} \{\chi_j(\mathbf{c}) | \mathbf{c} \in C^+\}.$$

Note that $X_j = \mathbf{1}$ for $j < 0$. For all integers i , define

$$Y_i \stackrel{\text{def}}{=} \{\chi_{-i-1}(\mathbf{c}) | \mathbf{c} \in C^-\}.$$

Note that $Y_i = \mathbf{1}$ for $i > 0$. Since the group trellis is time invariant, we have that $X_j \subset B$ for all j , and $Y_i \subset B$ for all i . Note that $X_j^+ = X_{j+1}^-$ and $Y_i^+ = Y_{i-1}^-$ for all integers i, j . And $X_j \triangleleft B$ and $Y_i \triangleleft B$ for all integers i, j [4].

The trellis states at time epoch t are Σ_t . A branch b_t at time epoch t has left state in Σ_t and right state in Σ_{t+1} . For any integer $l > 0$, we say a group trellis C is *l -controllable* if for any time epoch t , and any pair of states s and s' , where $s \in \Sigma_t$ and $s' \in \Sigma_{t+l}$, there is a trellis path through these two states. This means there is a trellis path segment of length l connecting the two states. The least integer l for which a group trellis is *l -controllable* is denoted as ℓ . In this paper, we only study the case $l = \ell$.

The next result follows directly from Proposition 7.2 of [4], using our notation.

Proposition 2 *The group trellis C is ℓ -controllable if and only if $(X_{\ell-1})^+ = B^-$, or equivalently, if and only if $(Y_{\ell-1})^- = B^+$.*

For each branch $b \in B$, we define the *next branch set* $\mathcal{N}(b)$ to be the set of branches that can follow b at the next time epoch in valid trellis paths. In other words, branch $e \in \mathcal{N}(b)$ if and only if $b^+ = e^-$.

Then the next branch set $\mathcal{N} : B \rightarrow B$ represents the contraction, equivalence, and expansion:

$$\frac{B}{Y_0} \simeq B^+ = \Sigma = B^- \simeq \frac{B}{X_0},$$

and more specifically for a branch $b \in B$, the assignments:

$$b \mapsto b^+ = s = e^- \mapsto X_0 e.$$

Proposition 3 *If $b^+ = e^-$, the next branch set $\mathcal{N}(b)$ of a branch b in B is the coset $X_0 e$ in B .*

For a set $H \subset B$, define the set $\mathcal{N}(H)$ to be the union $\cup_{b \in H} \mathcal{N}(b)$. The set $\mathcal{N}(H)$ always consists of cosets of X_0 . Note that $\mathcal{N}(X_j) = X_{j+1}$ for all integers $j \geq -1$.

Proposition 4 *If $b^+ = e^-$, the next branch set $\mathcal{N}(b)$ of a branch b in X_j is the coset $X_0 e$ in X_{j+1} , for $j \geq -1$.*

For a set $H \subset B$ and integer $j > 0$, define $\mathcal{N}^j(H)$ to be the j -fold composition $\mathcal{N}^j(H) = \mathcal{N} \circ \mathcal{N} \circ \dots \circ \mathcal{N}(H)$. For $j = 0$, define $\mathcal{N}^j(H) = \mathcal{N}^0(H)$ to be just H . If H is a set of trellis branches at time epoch 0, then $\mathcal{N}^j(H)$ is the set of trellis branches at time epoch j , such that for each $b_j \in \mathcal{N}^j(H)$ there is a $b_0 \in H$ and a path in the trellis from b_0 to b_j . Note that $X_j = \mathcal{N}^j(X_0)$.

For a set $H \subset B$ and integer $k \geq 0$, define $\mathcal{N}^{[0, k]}(H)$ to be the set of all trellis path segments $(b_0, \dots, b_k)$ on time interval $[0, k]$ that start with a branch $b_0 \in H$.

Proposition 5 *If G, H are subsets of B , then $\mathcal{N}(GH) = \mathcal{N}(G)\mathcal{N}(H)$.*

Proof. Let (g, g') be a trellis path section of length 2, with $g \in G$, and similarly for (h, h') . Then $(g, g') * (h, h') = (gh, g'h')$ is a trellis path section of length 2, with $gh \in GH$. Then $\mathcal{N}(gh) = \mathcal{N}(g)\mathcal{N}(h)$, so $\mathcal{N}(GH) = \mathcal{N}(G)\mathcal{N}(H)$. •

This result means $\mathcal{N}(Gb) = \mathcal{N}(G)\mathcal{N}(b)$ and $\mathcal{N}(bG) = \mathcal{N}(b)\mathcal{N}(G)$ for right coset Gb and left coset bG .

Proposition 6 *For any subsets G, H of B , we have $(G \cap H)^+ = G^+ \cap H^+$ and $(G \cap H)^- = G^- \cap H^-$. Also $(GH)^+ = G^+H^+$ and $(GH)^- = G^-H^-$.*

3. SCHREIER SERIES AND CONTROLLABLE MATRIX

The group B has two normal series (and chief series)

$$1 = X_{-1} \subset X_0 \subset X_1 \subset \cdots \subset X_\ell = B$$

and

$$1 = Y_{-1} \subset Y_0 \subset Y_1 \subset \cdots \subset Y_\ell = B.$$

We denote these normal series by $\{X_j\}$ and $\{Y_i\}$. The Schreier refinement theorem used to prove the Jordan-Hölder theorem [10] shows how to obtain a refinement of $\{X_j\}$ by inserting $\{Y_i\}$; we call this the *Schreier series* of $\{X_j\}$ and $\{Y_i\}$. Since $\{X_j\}$ and $\{Y_i\}$ are chief series, the Schreier series of $\{X_j\}$ and $\{Y_i\}$ is a chief series. In equation (1) (see next page), we have written the Schreier series as a matrix of $\ell+1$ columns and $\ell+2$ rows. Note that the terms in the bottom row form the sequence $X_{-1}, X_0, X_1, \dots, X_{\ell-2}, X_{\ell-1}$, and the terms in the top row form the sequence $X_0, X_1, X_2, \dots, X_{\ell-1}, X_\ell$. Thus (1) is indeed a refinement of the normal series $\{X_j\}$. We call (1) the *matrix form* of the Schreier series of $\{X_j\}$ and $\{Y_i\}$, or just the matrix form of $\{X_j\}$ and $\{Y_i\}$.

The normal chains $\{X_j\}$ and $\{Y_i\}$ were first studied in [4]. Factors of the diagonal terms, $X_j \cap Y_{\ell-j}$, for $0 \leq j \leq \ell$, are the groups used in [4] to give an abstract characterization of the branch group of an ℓ -controllable group trellis.

Pick the term in the upper left corner of the matrix form, $X_{-1}(X_0 \cap Y_\ell)$, and assume the group trellis is ℓ -controllable. Then

$$X_{-1}(X_0 \cap Y_\ell) = X_0.$$

Applying $\mathcal{N}$ to both sides, we have $\mathcal{N}(X_{-1}(X_0 \cap Y_\ell)) = X_0(X_1 \cap Y_{\ell-1})$ and $\mathcal{N}(X_0) = X_1$. Then

$$X_0(X_1 \cap Y_{\ell-1}) = X_1.$$

Continuing in this way we obtain the following.

Proposition 7 *If the group trellis C is ℓ -controllable, then*

$$X_{j-1}(X_j \cap Y_{\ell-j}) = X_j,$$

for $j \geq 0$.

Proposition 7 is essentially Proposition 7.2 of [4], obtained by a slightly different method.

The *diagonal terms* of the matrix form are $X_{j-1}(X_j \cap Y_{\ell-j})$ for $j = 0, \dots, \ell$. Proposition 7 shows that the diagonal terms satisfy $X_{j-1}(X_j \cap Y_{\ell-j}) = X_j$ for $j = 0, \dots, \ell$, if the group trellis is ℓ -controllable. For $j \in [1, \ell]$, this means all column terms above the diagonal term are the same as the diagonal term. Then we can reduce the matrix form to a triangular form as shown in (2). A triangle can be formed in two ways, depending on whether the columns in (1) are shifted up or not; we have shifted the columns up since it is more useful here. We call (2) the *controllable matrix* of $\{X_j\}$ and $\{Y_i\}$.

Theorem 8 *The Schreier series is a description (normal chain and chief series) of the branch group of a group trellis. The controllable matrix is a description (normal chain and chief series) of the branch group of a controllable group trellis.*

Proof. Both $\{X_j\}$ and $\{Y_i\}$ are normal chains of the branch group B . Then by the Schreier refinement theorem, the Schreier refinement is a normal chain of the branch group B . •

The *Schreier series* of $\{Y_i\}$ and $\{X_j\}$ is obtained by interchanging X and Y in (1); it is the dual of the Schreier series of $\{X_j\}$ and $\{Y_i\}$. In similar fashion, the *controllable matrix* of $\{Y_i\}$ and $\{X_j\}$ is the dual of the controllable matrix of $\{X_j\}$ and $\{Y_i\}$.

4. STRUCTURE OF THE CONTROLLABLE MATRIX

We index the rows and columns of (2), and denote terms, in a definite way. We index the columns with j , for $0 \leq j \leq \ell$, and rows with k , for $0 \leq k \leq \ell$, counting up from the bottom (in general, we do not include terms of the form $X_k(1)$). For example, the term $X_{-1}(X_0 \cap Y_0)$ is in column $j = 0$, and row $k = 0$. In general, we indicate a term in the controllable matrix by $X_{j-1}(X_j \cap Y_{k-j})$, where the subscripts mean definite things. The subscript α of X in the factor term $(X_\alpha \cap Y_\beta)$ always indicates the column, and the sum of the subscripts $\alpha + \beta$ of X and Y in the factor term always indicates the row. So the term $X_{j-1}(X_j \cap Y_{k-j})$ is in column j and row k . As other examples, the factor term $(X_j \cap Y_{k-j-1})$ is in column j and row $k - 1$, and the factor term $(X_{j-1} \cap Y_{k-j})$ is in column $j - 1$ and row $k - 1$. Note that row k of the controllable matrix has (length) $k + 1$ terms, ignoring the last term $X_k(1)$.

We now show the controllable matrix preserves shifts, that is, it has a **shift property**.

Proposition 9 *Fix k , $0 \leq k \leq \ell$, and fix j , $0 \leq j \leq k$. The controllable matrix has a shift property: the term $X_j(X_{j+1} \cap Y_{k-j-1})$ in column $j + 1$ and row k is a shift of the term $X_{j-1}(X_j \cap Y_{k-j})$ in column j and row k , that is*

$$\mathcal{N}(X_{j-1}(X_j \cap Y_{k-j})) = X_j(X_{j+1} \cap Y_{k-j-1}). \quad (3)$$

$$\begin{array}{cccccc}
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_\ell) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_\ell) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_\ell) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_\ell) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_\ell) \end{array} \\
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_{\ell-1}) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_{\ell-1}) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_{\ell-1}) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_{\ell-1}) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_{\ell-1}) \end{array} \\
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_{\ell-2}) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_{\ell-2}) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_{\ell-2}) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_{\ell-2}) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_{\ell-2}) \end{array} \\
\begin{array}{c} \cup \\ \cdots \end{array} & \begin{array}{c} \cup \\ \cdots \end{array} & \begin{array}{c} \cup \\ \cdots \end{array} & \cdots & \begin{array}{c} \cup \\ \cdots \end{array} & \begin{array}{c} \cup \\ \cdots \end{array} \\
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_2) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_2) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_2) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_2) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_2) \end{array} \\
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_1) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_1) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_1) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_1) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_1) \end{array} \\
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_0) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_0) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_0) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_0) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_0) \end{array} \\
\begin{array}{c} \cup \\ X_{-1}(X_0 \cap Y_{-1}) \end{array} & \begin{array}{c} \cup \\ X_0(X_1 \cap Y_{-1}) \end{array} & \begin{array}{c} \cup \\ X_1(X_2 \cap Y_{-1}) \end{array} & \cdots & \begin{array}{c} \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_{-1}) \end{array} & \begin{array}{c} \cup \\ X_{\ell-1}(X_\ell \cap Y_{-1}) \end{array}
\end{array} \tag{1}$$

$$\begin{array}{ccccccc}
\begin{array}{c} \parallel \\ X_{-1}(X_0 \cap Y_\ell) \\ \cup \\ X_{-1}(X_0 \cap Y_{\ell-1}) \\ \cup \\ X_{-1}(X_0 \cap Y_{\ell-2}) \\ \cup \\ \dots \\ \cup \\ X_{-1}(X_0 \cap Y_2) \\ \cup \\ X_{-1}(X_0 \cap Y_1) \\ \cup \\ X_{-1}(X_0 \cap Y_0) \\ \cup \\ X_{-1}(\mathbf{1}) \end{array} &
\begin{array}{c} \parallel \\ X_0(X_1 \cap Y_{\ell-1}) \\ \cup \\ X_0(X_1 \cap Y_{\ell-2}) \\ \cup \\ X_0(X_1 \cap Y_{\ell-3}) \\ \cup \\ \dots \\ \cup \\ X_0(X_1 \cap Y_1) \\ \cup \\ X_0(X_1 \cap Y_0) \\ \cup \\ X_0(\mathbf{1}) \end{array} &
\begin{array}{c} \parallel \\ X_1(X_2 \cap Y_{\ell-2}) \\ \cup \\ X_1(X_2 \cap Y_{\ell-3}) \\ \cup \\ X_1(X_2 \cap Y_{\ell-4}) \\ \cup \\ \dots \\ \cup \\ X_1(X_2 \cap Y_0) \\ \cup \\ X_1(\mathbf{1}) \end{array} &
\cdots &
\begin{array}{c} \parallel \\ X_{\ell-2}(X_{\ell-1} \cap Y_1) \\ \cup \\ X_{\ell-2}(X_{\ell-1} \cap Y_0) \\ \cup \\ X_{\ell-2}(\mathbf{1}) \end{array} &
\begin{array}{c} \parallel \\ X_{\ell-1}(X_\ell \cap Y_0) \\ \cup \\ X_{\ell-1}(\mathbf{1}) \end{array} &
X_\ell(\mathbf{1})
\end{array} \tag{2}$$

Proof. Fix k , $0 \leq k \leq \ell$, and fix j , $0 \leq j \leq k$. We have

$$(X_j \cap Y_{k-j})^+ = (X_{j+1} \cap Y_{k-j-1})^-, \quad (4)$$

By definition, we have $X_j^+ = X_{j+1}^-$ and $Y_{k-j}^+ = Y_{k-j-1}^-$. Then (4) follows from Proposition 6. The result (4) was previously given in [4].

Using (4), we have $\mathcal{N}(X_j \cap Y_{k-j}) = X_0(X_{j+1} \cap Y_{k-j-1})$. Then (3) follows using Proposition 5. •

The first column of the controllable matrix (2) will be important to us so we define $\Delta_k = X_0 \cap Y_k$, for $0 \leq k \leq \ell$; define $\Delta_{-1} = X_0 \cap Y_{-1} = \mathbf{1}$. We now show that row k of the controllable matrix is just $\mathcal{N}^{[0,k]}(\Delta_k)$, or just the path segments in C^+ on time interval $[0, k]$ that start with a branch $b_0 \in \Delta_k$.

Theorem 10 Fix k , $0 \leq k \leq \ell$. We have

$$\mathcal{N}^j(\Delta_k) = X_{j-1}(X_j \cap Y_{k-j}), \quad (5)$$

for $0 \leq j \leq k$. And $\mathcal{N}^j(\Delta_k) = X_{j-1}$ for $k < j \leq \ell$, $\mathcal{N}^j(\Delta_k) = X_\ell$ for $j > \ell$. The k^{th} row of the controllable matrix is just the terms in $\mathcal{N}^{[0,k]}(\Delta_k)$.

Proof. We prove (5) by induction. Assume it is true for $j = n$. Then use (3) to show it is true for $j = n + 1$. Then (5) shows the k^{th} row of the controllable matrix is just the terms in $\mathcal{N}^{[0,k]}(\Delta_k)$. •

Using Theorem 10, we can represent a quotient group of terms in the same column of (2) and adjacent rows in two different ways:

$$\frac{\mathcal{N}^j(\Delta_k)}{\mathcal{N}^j(\Delta_{k-1})} = \frac{X_{j-1}(X_j \cap Y_{k-j})}{X_{j-1}(X_j \cap Y_{k-j-1})}, \quad (6)$$

for $0 \leq j \leq k$.

We now show that terms in the controllable matrix in (2) can be related to certain paths in the trellis. Consider the portion of the trellis defined by the following projection:

$$P_{[0,\ell]} \stackrel{\text{def}}{=} \chi_{[0,\ell]}(C_{[0,\infty)}).$$

We call this portion of the trellis, $P_{[0,\ell]}$, the *pletty*. These are paths in the trellis which split from the identity state at time epoch 0. Note that $\text{pletty } P_{[0,\ell]} = \mathcal{N}^{[0,\ell]}(X_0)$. The branches in these paths are X_0 at time epoch 0, X_1 at time epoch 1, ..., and X_ℓ at time epoch ℓ . The branches at time epoch j in the pletty are X_j , for $0 \leq j \leq \ell$. We call X_j the j^{th} *pletty section*. Thus a pletty consists of the pletty sections $X_0, X_1, \dots, X_\ell$.

Proposition 11 Any pletty is a group, called the group pletty.

Proof. The projection map is a homomorphism, so the image is a group [3]. •

Proposition 12 Any pletty section is a group.

Proof. X_j is a group. •

The elements of $X_j = X_{j-1}(X_j \cap Y_{\ell-j})$ form a subgraph of the group trellis section B , where each input state at time j has $|X_0|$ splitting branches and each output state at time $j + 1$ has $|X_j \cap Y_0|$ merging branches. The right states of X_j , X_j^+ , are isomorphic to cosets of $X_j \cap Y_0$. The left states of X_j , X_j^- , are isomorphic to cosets of X_0 . Then X_j is a subdirect product of $X_j^- \times X_j^+$, a subgroup of the subdirect product B .

Of course we know the right states of X_j , X_j^+ , are connected to the left states of X_{j+1} , X_{j+1}^- , in the trellis. Thus the pletty sections connect to form the pletty.

It can also be seen that column j of (2), $0 \leq j \leq \ell$, represents the coset decomposition of the quotient group X_j/X_{j-1} . The right states of branches in column j are the left states of branches in column $j + 1$, $0 \leq j < \ell$. And we have just seen in Theorem 10 that the k^{th} row of the controllable matrix is just the path segments in the pletty that split from Δ_k . Thus (2) gives a “picture” of the groups that occur in the group pletty.

We can summarize the preceding discussion with the following theorem, which can be compared with Theorem 8.

Theorem 13 The controllable matrix is a description of the group pletty of a controllable group trellis.

Proposition 14 $\mathcal{N}^{[0,k]}(\Delta_{k-1})$ and $\mathcal{N}^{[0,k]}(\Delta_k)$ are groups.

Proof. $\mathcal{N}^{[0,k]}(\Delta_{k-1})$ is the projection of all paths in C^+ that have first nontrivial component in Δ_{k-1} on the interval $[0, k]$. •

Proposition 15 $\mathcal{N}^{[0,k]}(\Delta_{k-1}) \triangleleft \mathcal{N}^{[0,k]}(\Delta_k)$ if and only if $\Delta_{k-1} \triangleleft \Delta_k$.

Theorem 16 We have

$$\frac{\mathcal{N}^{[0,k]}(\Delta_k)}{\mathcal{N}^{[0,k]}(\Delta_{k-1})} \simeq \frac{\Delta_k}{\Delta_{k-1}}.$$

Proof. The projection $\chi_0 : \mathcal{N}^{[0,k]}(\Delta_k) \rightarrow \Delta_k$ is onto. It is a homomorphism with kernel $\mathcal{N}^{[0,k]}(\mathbf{1})$. The projection $\chi_0 : \mathcal{N}^{[0,k]}(\Delta_{k-1}) \rightarrow \Delta_{k-1}$ is onto. It is a homomorphism with kernel $\mathcal{N}^{[0,k]}(\mathbf{1})$. Therefore, by the first homomorphism theorem,

$$\begin{aligned} \frac{\mathcal{N}^{[0,k]}(\Delta_k)}{\mathcal{N}^{[0,k]}(\mathbf{1})} &\simeq \Delta_k, \\ \frac{\mathcal{N}^{[0,k]}(\Delta_{k-1})}{\mathcal{N}^{[0,k]}(\mathbf{1})} &\simeq \Delta_{k-1}. \end{aligned}$$

Now use the correspondence theorem and third isomorphism theorem to complete the proof. •

Proposition 17 $\mathcal{N}^j(\Delta_{k-1})$ and $\mathcal{N}^j(\Delta_k)$ are groups.

Proof. $\mathcal{N}^j(\Delta_{k-1})$ is the projection of all paths in C^+ that have first nontrivial component in Δ_{k-1} , or the projection of $\mathcal{N}^{[0,k]}(\Delta_{k-1})$, on the interval $[j, j]$. Alternatively, see (5).

Proposition 18 $\mathcal{N}^j(\Delta_{k-1}) \triangleleft \mathcal{N}^j(\Delta_k)$.

Proof. See (6).

Theorem 19 For $0 \leq j \leq k$, we have

$$\frac{\mathcal{N}^{[0,k]}(\Delta_k)}{\mathcal{N}^{[0,k]}(\Delta_{k-1})} \simeq \frac{\mathcal{N}^j(\Delta_k)}{\mathcal{N}^j(\Delta_{k-1})}.$$

Proof. The projection $\chi_j : \mathcal{N}^{[0,k]}(\Delta_k) \rightarrow \mathcal{N}^j(\Delta_k)$ is onto. It is a homomorphism with kernel K_k , the path segments in $\mathcal{N}^{[0,k]}(\Delta_k)$ that are the identity at time j . The projection $\chi_j : \mathcal{N}^{[0,k]}(\Delta_{k-1}) \rightarrow \mathcal{N}^j(\Delta_{k-1})$ is onto. It is a homomorphism with kernel K_{k-1} , the path segments in $\mathcal{N}^{[0,k]}(\Delta_{k-1})$ that are the identity at time j . Therefore, by the first homomorphism theorem,

$$\frac{\mathcal{N}^{[0,k]}(\Delta_k)}{K_k} \simeq \mathcal{N}^j(\Delta_k),$$

$$\frac{\mathcal{N}^{[0,k]}(\Delta_{k-1})}{K_{k-1}} \simeq \mathcal{N}^j(\Delta_{k-1}).$$

But $K_k = K_{k-1}$ since the path segments that are the identity at time j are in $X_0 \cap Y_{j-1}$, and $X_0 \cap Y_{j-1} \subset \Delta_{k-1} \subset \Delta_k$. Now use the correspondence theorem and third isomorphism theorem to complete the proof. •

Define

$$\Lambda_{[0,k]} \stackrel{\text{def}}{=} \frac{\mathcal{N}^{[0,k]}(\Delta_k)}{\mathcal{N}^{[0,k]}(\Delta_{k-1})}.$$

For $j = 0, 1, \dots, k$, by the j^{th} components of a transversal, we mean the set of branches which are the time j branch in each of the coset representatives in the transversal.

Corollary 20 The j^{th} components of a transversal of $\Lambda_{[0,k]}$ are a transversal of

$$\frac{\mathcal{N}^j(\Delta_k)}{\mathcal{N}^j(\Delta_{k-1})}, \quad (7)$$

for $0 \leq j \leq k$.

Proof. Theorem 19 shows that the projection $\chi_j(\Lambda_{[0,k]})$ gives a 1-1 correspondence between cosets of $\Lambda_{[0,k]}$ and cosets of (7). Therefore the projection χ_j of a transversal of $\Lambda_{[0,k]}$ is a transversal of (7). •

Corollary 21 For $0 \leq k \leq \ell$, and $0 \leq j \leq k$, we have

$$\frac{\Delta_k}{\Delta_{k-1}} \simeq \frac{\mathcal{N}^{[0,k]}(\Delta_k)}{\mathcal{N}^{[0,k]}(\Delta_{k-1})} \simeq \frac{\mathcal{N}^j(\Delta_k)}{\mathcal{N}^j(\Delta_{k-1})}.$$

Remark: This result can be regarded as a rectangle criterion for a controllable matrix, with Δ_k , Δ_{k-1} , $\mathcal{N}^j(\Delta_k)$, and $\mathcal{N}^j(\Delta_{k-1})$ as the corners of a rectangle in (2). It is similar in spirit to a quadrangle criterion for a Latin square [13] or a configuration theorem for a net [14]. In fact, the rectangle condition can be generalized further by starting with groups Δ_k and Δ_{k-m} , for $m > 1$. These more general results are not needed.

Corollary 21 shows the controllable matrix preserves quotient groups. For $0 \leq k \leq \ell$, and $0 \leq j \leq k$, define quotient group

$$Q_{j,k} \stackrel{\text{def}}{=} \mathcal{N}^j(\Delta_k) / \mathcal{N}^j(\Delta_{k-1}).$$

The $Q_{j,k}$ are the factor groups of the normal chain given by the controllable matrix, as shown in (16) on page 10. Corollary 21 gives $Q_{0,k} \simeq Q_{1,k} \simeq \dots \simeq Q_{j,k} \simeq \dots \simeq Q_{k,k}$ for $0 \leq k \leq \ell$. We use this result to obtain a natural shift structure of the controllable matrix in Section 6.

We now show the Forney-Trott generators are a transversal of $\Lambda_{[0,k]}$, and therefore components of the generators are a transversal of $Q_{j,k}$, $0 \leq j \leq k$. Forney and Trott [3] define a generator for a group code $\mathcal{C}$ using the quotient group

$$\frac{\mathcal{C}_{[0,k]}}{\mathcal{C}_{[0,k]}\mathcal{C}_{(0,k]}},$$

for $0 \leq k \leq \ell$. We can transcribe their approach to the group trellis C , used here, as

$$\Gamma_{[0,k]} \stackrel{\text{def}}{=} \frac{C_{[0,k]}}{C_{[0,k]}C_{(0,k]}},$$

where quotient group $\Gamma_{[0,k]}$ is called the *granule* [3]. Let $[\Gamma_{[0,k]}]$ be a set of coset representatives of $\Gamma_{[0,k]}$. We always take the coset representative of $C_{[0,k]}C_{(0,k]}$ to be the identity sequence. In case $\Gamma_{[0,k]}$ is isomorphic to the identity group, the identity sequence is the only coset representative. We call the coset representatives of $\Gamma_{[0,k]}$ *generators* $\mathbf{g}_{0,k}$. A nonidentity generator $\mathbf{g}_{0,k}$ is an element of $C_{[0,k]}$ but not of $C_{[0,k]}$ or of $C_{(0,k]}$, so its span is exactly $k+1$. Thus every nonidentity generator $\mathbf{g}_{0,k}$ is a codeword that cannot be expressed as a combination of shorter codewords [3].

For two sets $H_1, H_2 \subset B$ such that $H_1^+ = H_2^-$, define the *concatenation* of H_1 and H_2 , $H_1 \wedge H_2$, to be all the (valid) trellis path segments of length two with first component in H_1 and second component in H_2 .

Lemma 22 The set of paths formed by the concatenation of groups

$$\dots, \mathbf{1}, \mathbf{1} \wedge (X_0 \cap Y_k) \wedge \dots \wedge (X_j \cap Y_{k-j}) \wedge (X_{j+1} \cap Y_{k-j-1}) \wedge \dots \wedge (X_k \cap Y_0) \wedge \mathbf{1}, \mathbf{1}, \dots \quad (8)$$

is $C_{[0,k]}$.

Proof. From (4), we know

$$(X_j \cap Y_{k-j})^+ = (X_{j+1} \cap Y_{k-j-1})^-.$$

This means the set of paths formed by the concatenation of groups in (8) is well defined: for any branch $b \in X_j \cap Y_{k-j}$, there is a branch $b' \in X_{j+1} \cap Y_{k-j-1}$ such that $b^+ = b^-$, and (b, b') is part of a path. The paths consist of code sequences which split from the identity state at time epoch $t \geq 0$ and merge to the identity state at time epochs $t \leq k+1$.

Fix integer k such that $0 \leq k \leq \ell$. Let $\mathbf{c}$ be a sequence in $C_{[0,k]}$. At time epoch j , $0 \leq j \leq k$, c_j must be in X_j , but cannot be in X_m , $m > j$. Similarly, c_j must be in Y_{k-j} . Then $c_j \in X_j \cap Y_{k-j}$ for all $j \in [0, k]$. Since (8) contains all code sequences whose component $c_j \in X_j \cap Y_{k-j}$ for all $j \in [0, k]$, then $\mathbf{c}$ is in (8). Conversely, any path in (8) must be in $C_{[0,k]}$.

Lemma 23 For j , $0 \leq j \leq k$, we have $\chi_j(C_{[0,k]}) = X_j \cap Y_{k-j}$. For example, this means $X_0 \cap Y_k = \chi_0(C_{[0,k]})$ and $Y_0 \cap X_k = \chi_k(C_{[0,k]})$.

Proof. From (8), we know $\chi_j(C_{[0,k]}) \subset X_j \cap Y_{k-j}$.

For any $b \in X_j \cap Y_{k-j}$, there is a sequence $\mathbf{c} \in C_{[0,k]}$ such that $c_j = b$. Then we know $X_j \cap Y_{k-j} \subset \chi_j(C_{[0,k]})$. Then $\chi_j(C_{[0,k]}) = X_j \cap Y_{k-j}$.

Lemma 24 We have

$$\chi_{[0,k]}(C_{[0,k]}) \subset \mathcal{N}^{[0,k]}(\Delta_k), \quad (9)$$

and

$$\chi_{[0,k]}(C_{[0,k]}C_{(0,k]}) \subset \mathcal{N}^{[0,k]}(\Delta_{k-1}). \quad (10)$$

Proof. We have (9) holds if and only if $\chi_0(C_{[0,k]}) \subset \Delta_k$. But this follows from Lemma 23. We have (10) holds if and only if $\chi_0(C_{[0,k]}C_{(0,k]}) \subset \Delta_{k-1}$. But $\chi_0(C_{[0,k]}C_{(0,k]}) = \chi_0(C_{[0,k]}) = \Delta_{k-1}$.

Theorem 25 There is a one to one correspondence μ between cosets of $\Gamma_{[0,k]}$ and $\Lambda_{[0,k]}$, given by

$$\mu : C_{[0,k]}C_{(0,k]}\mathbf{c} \mapsto \mathcal{N}^{[0,k]}(\Delta_{k-1}c_0), \quad (11)$$

for $\mathbf{c} \in C_{[0,k]}$, where $\chi_0(C_{[0,k]}C_{(0,k]}\mathbf{c}) = \Delta_{k-1}c_0$, and $c_0 = \chi_0(\mathbf{c})$, such that

$$\chi_{[0,k]}(C_{[0,k]}C_{(0,k]}\mathbf{c}) \subset \mathcal{N}^{[0,k]}(\Delta_{k-1}c_0), \quad (12)$$

and

$$\Gamma_{[0,k]} \stackrel{\mu}{\cong} \Lambda_{[0,k]}.$$

Proof. We have that (12) holds if and only if $\chi_0(C_{[0,k]}C_{(0,k]}\mathbf{c}) \subset \Delta_{k-1}c_0$. But

$$\begin{aligned} \chi_0(C_{[0,k]}C_{(0,k]}\mathbf{c}) &= \chi_0(C_{[0,k]}\mathbf{c}) \\ &= \Delta_{k-1}c_0. \end{aligned}$$

This shows (12). Then we can properly define the 1-1 correspondence μ between cosets of $\Gamma_{[0,k]}$ and $\Lambda_{[0,k]}$ as given in (11).

Forney and Trott [3] define an *input chain* $F_0 \subset F_1 \subset \dots \subset F_\ell$ by the projection $F_k \stackrel{\text{def}}{=} \chi_0(C_{[0,k]})$ for

$k = 0, 1, \dots, \ell$. Using Lemma 23, this gives $F_k = \Delta_k$. In their Input Granule Theorem [3], Forney and Trott show that $\Gamma_{[0,k]} \simeq F_k/F_{k-1}$ for k such that $0 < k \leq \ell$. Then we have

$$\Gamma_{[0,k]} \simeq F_k/F_{k-1} = \Delta_k/\Delta_{k-1}.$$

Then using Theorem 16,

$$\Gamma_{[0,k]} \simeq \Delta_k/\Delta_{k-1} \simeq \Lambda_{[0,k]}.$$

Following the correspondences given in the Input Granule Theorem of [3] and Theorem 16 shows that the isomorphism $\Gamma_{[0,k]} \simeq \Lambda_{[0,k]}$ is given by μ . •

Corollary 26 Let $[\Gamma_{[0,k]}]$ be a set of generators which is a transversal of $\Gamma_{[0,k]}$. Then $\{\chi_{[0,k]}(\mathbf{g}_{0,k}) | \mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]\}$ is a transversal of $\Lambda_{[0,k]}$.

Fix k such that $0 \leq k \leq \ell$. Let generator $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$,

$$\mathbf{g}_{0,k} = \dots, \mathbf{1}, \mathbf{1}, r_{0,k}, r_{1,k}, \dots, r_{j,k}, \dots, r_{k,k}, \mathbf{1}, \mathbf{1}, \dots, \quad (13)$$

be a representative in a transversal of $\Gamma_{[0,k]}$. From (8) we know component $r_{j,k}$ is an element of $X_j \cap Y_{k-j}$, and from Corollaries 26 and 20 we know $r_{j,k}$ is a representative of

$$Q_{j,k} = \frac{\mathcal{N}^j(\Delta_k)}{\mathcal{N}^j(\Delta_{k-1})} = \frac{X_{j-1}(X_j \cap Y_{k-j})}{X_{j-1}(X_j \cap Y_{k-j-1})}, \quad (14)$$

for $j = 0, 1, \dots, k$. Thus the transversal $[\Gamma_{[0,k]}]$ gives a set of transversals

$$[r_{0,k}], [r_{1,k}], \dots, [r_{j,k}], \dots, [r_{k,k}],$$

where $[r_{j,k}]$ is a transversal of (14), for $j = 0, 1, \dots, k$. Since this is true for each k , $0 \leq k \leq \ell$, we can obtain the matrix shown in (17) on page 10. It can be seen that array (17) corresponds to the controllable matrix (2) and $[r_{j,k}]$ corresponds to group elements in the controllable matrix. The controllable matrix gives a normal chain of B , and the group elements of (17) form a *complete system of coset representatives* of the normal chain. Thus any branch $b \in B$, and any $b_t \in B$, can be written using elements of (17) as

$$\begin{aligned} b_t &= r_{0,0}r_{0,1}r_{0,2} \dots r_{0,\ell}r_{1,1} \dots r_{1,\ell}r_{2,2} \dots \\ &\quad \dots r_{j,j} \dots r_{j,k} \dots r_{j,\ell} \dots r_{\ell,\ell}, \end{aligned} \quad (15)$$

where the terms in product $(r_{j,j} \dots r_{j,k} \dots r_{j,\ell})$ are coset representatives from the j -th column of (17), $0 \leq j \leq \ell$.

Theorem 27 The j^{th} components of generators $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$, for $0 \leq j \leq k$, for $0 \leq k \leq \ell$, form a complete system of coset representatives for the normal chain of B given by the controllable matrix of $\{X_j\}$ and $\{Y_i\}$.

The following theorem is interesting but not essential.

Theorem 28 The j^{th} component, $0 \leq j \leq k$, of a generator $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$ is an element of $X_j \cap Y_{k-j}$ taken from cosets of

$$\frac{(X_j \cap Y_{k-j})}{(X_j \cap Y_{k-j-1})(X_{j-1} \cap Y_{k-j})},$$

and in fact,

$$\frac{\mathcal{N}^j(\Delta_k)}{\mathcal{N}^j(\Delta_{k-1})} \simeq \frac{(X_j \cap Y_{k-j})}{(X_j \cap Y_{k-j-1})(X_{j-1} \cap Y_{k-j})}.$$

The theorem is easy to prove (see [16]) using (6) and the following lemma, excised from the proof of the Zassenhaus lemma (see p. 100 of [10]).

Lemma 29 (from proof of Zassenhaus lemma) Let $U \triangleleft U^*$ and $V \triangleleft V^*$ be four subgroups of a group G . Then $D = (U^* \cap V)(U \cap V^*)$ is a normal subgroup of $U^* \cap V^*$. If $g \in U(U^* \cap V^*)$, then $g = uu^*$ for $u \in U$ and $u^* \in U^* \cap V^*$. Define function $f : U(U^* \cap V^*) \rightarrow (U^* \cap V^*)/D$ by $f(g) = f(uu^*) = Du^*$. Then f is a well defined homomorphism with kernel $U(U^* \cap V)$ and

$$\frac{U(U^* \cap V^*)}{U(U^* \cap V)} \simeq \frac{U^* \cap V^*}{D}.$$

We can extend Theorem 27 to show that the components of generators $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$ form a complete system of coset representatives for two normal chains. Define $\Delta'_k = Y_0 \cap X_k$, for $0 \leq k \leq \ell$. Define the *previous branch set* $\mathcal{P}(b)$ to be the time reversal of $\mathcal{N}(b)$. Define quotient group

$$\Lambda'_{[0,k]} \stackrel{\text{def}}{=} \frac{\mathcal{P}^{[0,k]}(\Delta'_k)}{\mathcal{P}^{[0,k]}(\Delta'_{k-1})},$$

where $\mathcal{P}^{[0,k]}$ is the time reversal of $\mathcal{N}^{[0,k]}$. Using these definitions, the arguments in Lemma 24 and Theorem 25 can be reversed in time. In place of the input chain [3], the *last output chain* [3] is used. This gives the following time reversed version of Theorem 25.

Theorem 30 There is a one to one correspondence μ' between cosets of $\Gamma_{[0,k]}$ and $\Lambda'_{[0,k]}$, given by

$$\mu' : C_{[0,k]}C_{(0,k)}\mathbf{c} \mapsto \mathcal{P}^{[0,k]}(\Delta'_{k-1}c_k),$$

for $\mathbf{c} \in C_{[0,k]}$, where $\chi_k(C_{[0,k]}C_{(0,k)}\mathbf{c}) = \Delta'_{k-1}c_k$, and $c_k = \chi_k(\mathbf{c})$, such that

$$\chi_{[0,k]}(C_{[0,k]}C_{(0,k)}\mathbf{c}) \subset \mathcal{P}^{[0,k]}(\Delta'_{k-1}c_k),$$

and

$$\Gamma_{[0,k]} \stackrel{\mu'}{\simeq} \Lambda'_{[0,k]}.$$

Corollary 31 Let $[\Gamma_{[0,k]}]$ be a set of generators which is a transversal of $\Gamma_{[0,k]}$. Then $\{\chi_{[0,k]}(\mathbf{g}_{0,k}) | \mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]\}$ is a transversal of $\Lambda'_{[0,k]}$.

Combining Corollaries 26 and 31 gives the following extension of Theorem 27.

Theorem 32 The j^{th} components of generators $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$, for $0 \leq j \leq k$, for $0 \leq k \leq \ell$, form a complete system of coset representatives for two different normal chains, the normal chain of B given by the controllable matrix of $\{X_j\}$ and $\{Y_i\}$, and the normal chain of B given by the controllable matrix of $\{Y_i\}$ and $\{X_j\}$ (of course a different ordering of the coset representatives is used for the two normal chains).

For integer l such that $0 \leq l \leq \ell$, define $B^{(l)}$ to be the branch group of the l -controllable subcode [3], and $\{X_j^{(l)}\}$ and $\{Y_i^{(l)}\}$ to be normal chains of the subcode analogous to $\{X_j\}$ and $\{Y_i\}$. In a similar way we can show generator $\mathbf{g}_{0,k}$ can be used for the l -controllable subcodes.

Theorem 33 Fix l such that $0 \leq l \leq \ell$. The j^{th} components of generators $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$, for $0 \leq j \leq k$, for $0 \leq k \leq l$, form a complete system of coset representatives for two different normal chains, the normal chain of $B^{(l)}$ given by the controllable matrix of $\{X_j^{(l)}\}$ and $\{Y_i^{(l)}\}$, and the normal chain of $B^{(l)}$ given by the controllable matrix of $\{Y_i^{(l)}\}$ and $\{X_j^{(l)}\}$.

5. ENCODER FOR $\{X_j\}$ AND $\{Y_i\}$

An *encoder* of the group trellis is a finite state machine that, given a sequence of inputs, can produce any path (any sequence of states and branches) in the group trellis. An encoder can help to explain the structure of a group trellis. We give an encoder here which has a register structure and uses the same generators as in [3], but the encoder is different.

For each $\mathbf{g}_{0,k} \in [\Gamma_{[0,k]}]$, $0 \leq k \leq \ell$, and each time t , let $\mathbf{g}_{t,k}$ be the time shift of $\mathbf{g}_{0,k}$; $\mathbf{g}_{t,k}$ is the same as the identity path outside the time interval $[t, t+k]$. We say $\mathbf{g}_{t,k}$ is a *generator at time t* . Let $[\Gamma_{[t,t+k]}]$ be the set of generators $\mathbf{g}_{t,k}$, $0 \leq k \leq \ell$.

Fix time t . We now show how to realize any element $r_{j,k}$ in (15) at time t . First rewrite (15) as

$$b_t = \prod_{j=0}^{\ell} \left(\prod_{k=j}^{\ell} r_{j,k} \right). \quad (19)$$

For any element $r_{j,k}$ in (19), $0 \leq k \leq \ell$ and $0 \leq j \leq k$, there is a $\mathbf{g}_{t-j,k} \in [\Gamma_{[t-j,t-j+k]}]$ such that

$$\chi_t(\mathbf{g}_{t-j,k}) = r_{j,k}. \quad (20)$$

Therefore we can rewrite (19) as

$$b_t = \prod_{j=0}^{\ell} \left(\prod_{k=j}^{\ell} \chi_t(\mathbf{g}_{t-j,k}) \right). \quad (21)$$

Thus we have shown any element $b_t \in B_t$ can be written using generators selected at times in the interval $[t-\ell, t]$, i.e., generators

$$\mathbf{g}_{t-j,k}, \text{ for } j = 0, \dots, \ell, \text{ for } k = j, \dots, \ell.$$

$$\begin{array}{cccccccccc}
Q_{0,\ell} & Q_{1,\ell} & \cdots & \cdots & Q_{j,\ell} & \cdots & \cdots & \cdots & Q_{\ell-1,\ell} & Q_{\ell,\ell} \\
Q_{0,\ell-1} & Q_{1,\ell-1} & \cdots & \cdots & Q_{j,\ell-1} & \cdots & \cdots & \cdots & Q_{\ell-1,\ell-1} & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & \\
Q_{0,k} & Q_{1,k} & \cdots & \cdots & Q_{j,k} & \cdots & Q_{k,k} & & & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & & & \\
\cdots & \cdots & \cdots & \cdots & Q_{j,j} & & & & & \\
\vdots & \vdots & \vdots & & & & & & & \\
Q_{0,2} & Q_{1,2} & Q_{2,2} & & & & & & & \\
Q_{0,1} & Q_{1,1} & & & & & & & & \\
Q_{0,0} & & & & & & & & &
\end{array} \tag{16}$$

$$\begin{array}{cccccccccc}
[r_{0,\ell}] & [r_{1,\ell}] & \cdots & \cdots & [r_{j,\ell}] & \cdots & \cdots & \cdots & [r_{\ell-1,\ell}] & [r_{\ell,\ell}] \\
[r_{0,\ell-1}] & [r_{1,\ell-1}] & \cdots & \cdots & [r_{j,\ell-1}] & \cdots & \cdots & \cdots & [r_{\ell-1,\ell-1}] & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & \\
[r_{0,k}] & [r_{1,k}] & \cdots & \cdots & [r_{j,k}] & \cdots & [r_{k,k}] & & & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & & & \\
\cdots & \cdots & \cdots & \cdots & [r_{j,j}] & & & & & \\
\vdots & \vdots & \vdots & & & & & & & \\
[r_{0,2}] & [r_{1,2}] & [r_{2,2}] & & & & & & & \\
[r_{0,1}] & [r_{1,1}] & & & & & & & & \\
[r_{0,0}] & & & & & & & & &
\end{array} \tag{17}$$

$$\begin{array}{cccccccccc}
U_{0,\ell} & U_{1,\ell} & \cdots & \cdots & U_{j,\ell} & \cdots & \cdots & \cdots & U_{\ell-1,\ell} & U_{\ell,\ell} \\
U_{0,\ell-1} & U_{1,\ell-1} & \cdots & \cdots & U_{j,\ell-1} & \cdots & \cdots & \cdots & U_{\ell-1,\ell-1} & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & \\
U_{0,k} & U_{1,k} & \cdots & \cdots & U_{j,k} & \cdots & U_{k,k} & & & \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & & & \\
\cdots & \cdots & \cdots & \cdots & U_{j,j} & & & & & \\
\vdots & \vdots & \vdots & & & & & & & \\
U_{0,2} & U_{1,2} & U_{2,2} & & & & & & & \\
U_{0,1} & U_{1,1} & & & & & & & & \\
U_{0,0} & & & & & & & & &
\end{array} \tag{18}$$

We will define an encoder based on (21). We can rewrite b_t to separate out the $j = 0$ term as

$$b_t = \prod_{k=0}^{\ell} \chi_t(\mathbf{g}_{t,k}) \quad (22)$$

$$\prod_{j=1}^{\ell} \left(\prod_{k=j}^{\ell} \chi_t(\mathbf{g}_{t-j,k}) \right). \quad (23)$$

The term in (22) is just an element $x_t \in X_0$. Note that x_t is a function of generators at time t . We will think of x_t as an *input* of the encoder. The term in (23) is just a branch $\hat{b}_t \in B_t$; it corresponds to the branch obtained when $x_t = \mathbf{1}$. Note that $\hat{b}_t$ is a function of generators at times $t', t' < t$. We can rewrite b_t as $b_t = x_t \hat{b}_t$, where $\hat{b}_t \in B_t$ and

$$\hat{b}_t = \prod_{j=1}^{\ell} \left(\prod_{k=j}^{\ell} \chi_t(\mathbf{g}_{t-j,k}) \right).$$

Note that $\hat{b}_t$ can take on $|B/X_0|$ different values, and each value is in a distinct coset of B/X_0 . Thus there is a one to one correspondence between the set of $\hat{b}_t$, $\{\hat{b}_t\}$, and cosets in B/X_0 , $\{\hat{b}_t\} \leftrightarrow B/X_0$. Thus we can think of $\hat{b}_t$ as a *state* of the encoder.

We propose an encoder based on (21) which has the same form as (21) for each time epoch. Accordingly, the form of the encoder for time $t + 1$ is

$$b_{t+1} = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} \chi_{t+1}(\mathbf{g}'_{t+1-j,k}).$$

We can rewrite this to separate out the $j = 0$ term:

$$b_{t+1} = \prod_{k=0}^{\ell} \chi_{t+1}(\mathbf{g}'_{t+1,k}) \quad (24)$$

$$\prod_{j=1}^{\ell} \prod_{k=j}^{\ell} \chi_{t+1}(\mathbf{g}'_{t+1-j,k}). \quad (25)$$

The term in (24) uses $\ell + 1$ new generators at time $t + 1$, $\mathbf{g}'_{t+1,k}$, for $k = 0, \dots, \ell$. It is the new input $x_{t+1} \in X_0$. Let $\hat{b}_{t+1}$ be the term in (25). We see that $\hat{b}_{t+1}$ uses generators $\mathbf{g}'_{t+1-j,k}$ at times in the interval $[t - \ell + 1, t]$.

We complete the specification of the encoder by requiring that the generators in (25), or $\hat{b}_{t+1}$, used by the encoder at time $t + 1$ be the same as the generators used by the encoder at time t . In other words, we require

$$\mathbf{g}'_{t+1-j,k} = \mathbf{g}_{t+1-j,k}$$

for $j = 1, \dots, \ell$, for $k = j, \dots, \ell$. Then

$$\hat{b}_{t+1} = \prod_{j=1}^{\ell} \prod_{k=j}^{\ell} \chi_{t+1}(\mathbf{g}_{t+1-j,k}). \quad (26)$$

The encoder output at time $t + 1$ is $b_{t+1} = x_{t+1} \hat{b}_{t+1}$. Thus the encoder uses a sliding block encoding of the past, given by $\hat{b}_{t+1}$, with new inputs at each time epoch, given by x_{t+1} . With this definition of the encoder, note that for any input x_{t+1} to the encoder, (b_t, b_{t+1}) is a trellis path segment of length two. In particular, for $x_{t+1} = \mathbf{1}$, we have $(b_t, \hat{b}_{t+1})$ is a trellis path segment of length two.

We now verify that given an arbitrary path $\mathbf{c}^*$ in the group trellis, the encoder can track it perfectly, i.e., produce the same path. First we give a useful lemma. We think of the encoder as an estimator.

Lemma 34 *Let $\mathbf{c}$ and $\mathbf{c}^*$ be two paths in the group trellis C :*

$$\begin{aligned} \mathbf{c} &= \dots, b_t, b_{t+1}, b_{t+2}, \dots \\ \mathbf{c}^* &= \dots, b_t^*, b_{t+1}^*, b_{t+2}^*, \dots \end{aligned}$$

The two trellis paths are arbitrary except that at time t , $b_t = b_t^*$. Then at time $t + 1$, b_{t+1} is in the same coset of B/X_0 as b_{t+1}^* , i.e., $b_{t+1} \in X_0 b_{t+1}^*$. At time $t + 2$, b_{t+2} is in the same coset of B/X_1 as b_{t+2}^* . In general at time $t + j$, $j \leq \ell$, b_{t+j} is in the same coset of B/X_{j-1} as b_{t+j}^* . For time $t + j$, $j > \ell$, b_{t+j} is in the same coset of B/X_{ℓ} as b_{t+j}^* ; this coset is just B . In a similar way, at time $t - 1$, b_{t-1} is in the same coset of B/Y_0 as b_{t-1}^* .

Proof. We have $\mathcal{N}(b_t) = \mathcal{N}(b_t^*)$, so b_{t+1} and b_{t+1}^* are in the same coset of X_0 . The rest of the proof is analogous.

•

We can think of $\mathbf{c}$ as a path which estimates $\mathbf{c}^*$. The lemma shows how the estimation degrades as time goes on when it is perfect at $t = 0$, for two otherwise arbitrary paths. Note that the conclusion of the lemma is unchanged if, instead of one path $\mathbf{c}$, we use a finite number of paths to estimate $\mathbf{c}^*$.

We now show that the encoder can track arbitrary path $\mathbf{c}^*$. Fix time t . The branch in $\mathbf{c}^*$ at time t is b_t^* . First we show that the encoder can produce the initial condition b_t^* . But we already know the encoder can produce any branch $b_t = b_t^*$ (see (21)). Thus at time t , the encoder can produce the branch b_t^* in $\mathbf{c}^*$.

Now we show the encoder can track path $\mathbf{c}^*$. First we show the encoder can track $\mathbf{c}^*$ at time $t + 1$. First find $\hat{b}_{t+1}$ as given in (26). We know $(b_t, \hat{b}_{t+1})$ is a trellis path segment of length two. Therefore we know from Lemma 34 that $b_{t+1}^* \in X_0 \hat{b}_{t+1}$. Thus there exists $\hat{x}_{t+1} \in X_0$ such that $b_{t+1}^* = \hat{x}_{t+1} \hat{b}_{t+1}$. Now select $\ell + 1$ new generators at time $t + 1$, $\hat{\mathbf{g}}_{t+1,k}$, for $k = 0, \dots, \ell$, such that

$$\prod_{k=0}^{\ell} \chi_{t+1}(\hat{\mathbf{g}}_{t+1,k}) = \hat{x}_{t+1}.$$

We know this is possible since $\hat{x}_{t+1} \in X_0$ and the first column of (17) is a coset decomposition of X_0 . With input $\hat{x}_{t+1}$ at time $t + 1$, the encoder output is given by $\hat{x}_{t+1} \hat{b}_{t+1} = b_{t+1}^*$. Thus we see that with proper input

$\hat{x}_{t+1}$, the encoder can track path $\mathbf{c}^*$ at time $t+1$. We can repeat this same argument for succeeding time epochs.

We can observe several features of the encoder. The term in the parentheses of (21) is some function of $t-j$, say h_{t-j} . Then $b_t = \prod_{j=0}^{\ell} h_{t-j}$. Thus the encoder has the form of a time convolution, reminiscent of a linear system. The Forney-Trott encoder [3] and Loeliger-Mittelholzer encoder [4] do not have the convolution analogy.

Note that if we apply input $x_t = \chi_t(\mathbf{g}_{t,k})$ at time t to an encoder in the identity state, the output is just

$$\chi_{t+n}(\mathbf{g}_{t,k}), \text{ for } n = 0, 1, \dots, \infty.$$

Thus the output response of the encoder is the input generator. This is reminiscent of the impulse response of a linear system.

6. A NATURAL SHIFT STRUCTURE

6.1 The controllable matrix as a shift structure

In view of the controllable matrix (2), and (17), a natural interpretation of the encoder is shown in Figure 1. The encoder consists of $\ell + 1$ registers, where the j^{th} register, $j = 0, 1, \dots, \ell$, contains the information (coset representatives) in quotient group X_j/X_{j-1} . At the next time epoch, the register containing X_j/X_{j-1} is shifted or “shoved” into the register containing X_{j+1}/X_j . The interpretation of the registers shown in Figure 1 is somewhat different from the classic interpretation of a shift register.




Figure 1: Shift structure.

As time increases, we can imagine that the shift structure in Figure 1 slides along the controllable matrix from left to right in convolutional fashion. Then the registers of Figure 1 are successively filled with X_0 , X_1/X_0 , and so on.

We represent the shift structure as a bank of independent shift registers in Figure 2. There is a natural relationship between the shift register bank in Figure 2 and the quotient groups $Q_{j,k}$ of the controllable matrix in (16). For $0 \leq k \leq \ell$ and $0 \leq j \leq k$, define $U_{j,k}$ to be the set of integers $0, 1, \dots, |Q_{j,k}| - 1$. For $0 \leq j \leq k$, we associate an integer $u_{j,k} \in U_{j,k}$ with a coset $\mathcal{N}^j(\Delta_{k-1})g$

in quotient group $Q_{j,k}$, denoted by $\mathcal{N}^j(\Delta_{k-1})g \mapsto u_{j,k}$, in the following way: for $0 \leq j < k$, if $\mathcal{N}^j(\Delta_{k-1})g \mapsto u_{j,k}$ then $\mathcal{N}(\mathcal{N}^j(\Delta_{k-1})g) \mapsto u_{j+1,k}$, where $u_{j+1,k} = u_{j,k}$. This can be regarded as a shift map $\zeta_{j,k} : U_{j,k} \rightarrow U_{j+1,k}$, given by the assignment $\zeta_{j,k} : u_{j,k} \mapsto u_{j+1,k}$, which models a register row in Figure 2. Then we can represent the shift register bank in Figure 2 as a triangular matrix of $\ell + 1$ rows and $\ell + 1$ columns over the sets $U_{j,k}$, of the form shown in (18) on page 10. We let $\mathbf{U}$ be the set of all triangular $\ell + 1$ by $\ell + 1$ matrices of the form (18). Let $\mathbf{u}$ be an element of $\mathbf{U}$, that is a triangular $\ell + 1$ by $\ell + 1$ matrix.

Note that if Δ_k/Δ_{k-1} is trivial, i.e., $\Delta_k/\Delta_{k-1} = \mathbf{1}$, then the transversal of $\Lambda_{[0,k]} = \mathcal{N}^{[0,k]}(\Delta_k)/\mathcal{N}^{[0,k]}(\Delta_{k-1})$ is trivial and the generator representative $\mathbf{g}_{0,k}$ is selected to be the identity sequence. Then $Q_{j,k} = \mathbf{1}$ for $0 \leq j \leq k$, and $U_{j,k} = 0$ for $0 \leq j \leq k$. This means there is a trivial row of $k+1$ zeros in (18) and a trivial register row in Figure 2 always filled with $k+1$ zeros.




Figure 2: A shift register bank.

Let $\mathbf{U}_j$ be the set of all columns of $\mathbf{U}$, $0 \leq j \leq \ell$, and let $\mathbf{u}_j$ be an element of $\mathbf{U}_j$. Then $\mathbf{u} = (\mathbf{u}_0, \dots, \mathbf{u}_{\ell})$. A column $\mathbf{u}_j$ is an $\ell - j + 1$ by 1 vector of the form

$$\mathbf{u}_j = (u_{j,\ell} \cdots u_{j,k} \cdots u_{j,j})^T, \quad (27)$$

where each $u_{j,k}$, $j \leq k \leq \ell$, is an integer in $U_{j,k}$. We denote a column with all entries 0 by $\mathbf{0}$. We define a shift map $\sigma_j : \mathbf{U}_j \rightarrow \mathbf{U}_{j+1}$ on columns $\mathbf{U}_j$, $0 \leq j < \ell$, by the assignment

$$\sigma_j : \mathbf{u}_j \mapsto \mathbf{u}_{j+1},$$

where $\mathbf{u}_{j+1}$ is the $\ell - j$ by 1 vector with components

$$(u_{j+1,\ell} \cdots u_{j+1,k} \cdots u_{j+1,j+1})^T. \quad (28)$$

In other words, for $k > j$ each component $u_{j+1,k}$ of $\mathbf{u}_{j+1}$ is just a shift of component $u_{j,k}$ of $\mathbf{u}_j$, using $\zeta_{j,k} : u_{j,k} \mapsto u_{j+1,k}$ where $u_{j+1,k} = u_{j,k}$. Then it can be seen that $\mathbf{U}$ and shift σ_j model the shift register bank in Figure 2. We abbreviate $\sigma_j(\mathbf{u}_j)$ as $\sigma_j \mathbf{u}_j$.

In this paper, we only consider directed graphs. A *directed graph* $\mathcal{D}$ is defined by a set $\mathcal{V}(\mathcal{D})$ of elements called *nodes* (also *states*), a set $\mathcal{E}(\mathcal{D})$ of elements called *edges*, and an incidence relation between nodes and edges

which associates with each edge an ordered pair of nodes, called *endpoints*.

We now associate a graph $\mathcal{D}_U$ with U . Each triangular matrix $u \in U$ is associated with an edge in $\mathcal{D}_U$. The last ℓ columns of u are a triangular $\ell + 1$ by ℓ matrix u_s , associated with a node in $\mathcal{D}_U$. u_s is an element of U_s , the set of all triangular $\ell + 1$ by ℓ matrices formed by the last ℓ columns of U .

Lemma 35 *In $\mathcal{D}_U$, edge $u = (u_0, u_1, \dots, u_\ell)$ has endpoints (u_a, u_b) , where $u_a = (u_1, \dots, u_\ell)$, the last ℓ components of u , and $u_b = (\sigma_0 u_0, \sigma_1 u_1, \dots, \sigma_{\ell-1} u_{\ell-1})$, a shift of the first ℓ components of u .*

Then it can be seen that $\mathcal{D}_U$ is a graph which models the shift register bank in Figure 2.

6.2 The graph automorphism group of the shift register bank

Let $\mathcal{D}$ and $\mathcal{H}$ be directed graphs. Let φ_s be a 1-1 mapping of $\mathcal{V}(\mathcal{D})$ onto $\mathcal{V}(\mathcal{H})$, and φ_e a 1-1 mapping of $\mathcal{E}(\mathcal{D})$ onto $\mathcal{E}(\mathcal{H})$. Let φ denote the ordered pair (φ_s, φ_e) . We say that $\varphi = (\varphi_s, \varphi_e)$ is an *isomorphism of $\mathcal{D}$ onto $\mathcal{H}$* if the following condition holds: if edge e has endpoints (a, b) , then edge $\varphi_e(e)$ has endpoints $(\varphi_s(a), \varphi_s(b))$. Note that once φ_e is specified, then φ_s is determined and thus so is $\varphi = (\varphi_s, \varphi_e)$.

An *automorphism* of a graph $\mathcal{D}$ is an isomorphism of $\mathcal{D}$ onto $\mathcal{D}$. The set of automorphisms of a graph $\mathcal{D}$ forms an *automorphism group* $\text{Aut}(\mathcal{D})$, where the group operation is composition, and the composition of two automorphisms φ and φ' , $\varphi \circ \varphi'$, is formed as $\varphi \circ \varphi' = (\varphi_s \circ \varphi'_s, \varphi_e \circ \varphi'_e)$, where $\varphi = (\varphi_s, \varphi_e)$ and $\varphi' = (\varphi'_s, \varphi'_e)$.

We want to find the automorphism group of $\mathcal{D}_U$, $\text{Aut}(\mathcal{D}_U)$. If $\varphi = (\varphi_s, \varphi_e)$ is to be an automorphism of $\mathcal{D}_U$, we know that $\varphi_e : U \rightarrow U$ and $\varphi_s : U_s \rightarrow U_s$ must be 1-1 and onto. Define a component form of φ_e by $\varphi_e = (\varphi_{e,0}, \dots, \varphi_{e,\ell})$ where function $\varphi_{e,j} : U \rightarrow U_j$ gives the j^{th} component of φ_e , $j = 0, \dots, \ell$. Similarly, let $\varphi_s = (\varphi_{s,1}, \dots, \varphi_{s,\ell})$ where function $\varphi_{s,j} : U_s \rightarrow U_j$ gives the j^{th} component of φ_s , $j = 1, \dots, \ell$. Also, in order for φ to be an automorphism, we must have that if u has endpoints (u_a, u_b) , then $\varphi_e(u)$ has endpoints $(\varphi_s(u_a), \varphi_s(u_b))$. Let us use Lemma 35 to see what this implies about φ_e and φ_s . First, it means that the last ℓ components of $\varphi_e(u)$ are $\varphi_s(u_a)$, or

$$\varphi_{s,j}(u_1, \dots, u_\ell) = \varphi_{e,j}(u_0, u_1, \dots, u_\ell) \quad (29)$$

for $j = 1, \dots, \ell$. Second, it means that $\varphi_s(u_b)$ is a shift of the first ℓ components of $\varphi_e(u)$, or

$$\varphi_{s,j}(\sigma_0 u_0, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_{j-1} \varphi_{e,j-1}(u_0, u_1, \dots, u_\ell), \quad (30)$$

for $j = 1, \dots, \ell$.

We now examine (29) and (30) further. The left hand side of (29) is a function only of $u_1, \dots, u_\ell$, so $\varphi_{e,j}$ cannot

be a function of u_0 for $j = 1, \dots, \ell$. Thus we can rewrite (29) as

$$\varphi_{s,j}(u_1, \dots, u_\ell) = \varphi_{e,j}(-, u_1, \dots, u_\ell) \quad (31)$$

for $j = 1, \dots, \ell$, where a dash indicates $\varphi_{e,j}$ is not a function of that component. We can use this result in (30) to obtain the set of equations

$$\varphi_{e,1}(-, \sigma_0 u_0, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_0 \varphi_{e,0}(u_0, u_1, \dots, u_\ell), \quad (32)$$

and

$$\varphi_{e,j}(-, \sigma_0 u_0, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_{j-1} \varphi_{e,j-1}(-, u_1, \dots, u_\ell), \quad (33)$$

for $j = 2, \dots, \ell$.

We can reduce the set of equations (33) further. Start with $j = 2$,

$$\varphi_{e,2}(-, \sigma_0 u_0, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_1 \varphi_{e,1}(-, u_1, \dots, u_\ell).$$

Fix $u_1, \dots, u_\ell$ on the right hand side; then the value of the right hand side is fixed. Since $u_1, \dots, u_\ell$ are fixed on the right hand side, $\sigma_1 u_1, \dots, \sigma_{\ell-1} u_{\ell-1}$ are fixed on the left hand side. Then to have equality, $\varphi_{e,2}$ cannot be a function of $\sigma_0 u_0$, or

$$\varphi_{e,2}(-, -, \sigma_1 u_1, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_1 \varphi_{e,1}(-, u_1, \dots, u_\ell). \quad (34)$$

Now look at the case $j = 3$,

$$\varphi_{e,3}(-, \sigma_0 u_0, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_2 \varphi_{e,2}(-, u_1, \dots, u_\ell).$$

Using the result (34), we obtain

$$\varphi_{e,3}(-, \sigma_0 u_0, \dots, \sigma_{\ell-1} u_{\ell-1}) = \sigma_2 \varphi_{e,2}(-, -, u_2, \dots, u_\ell).$$

Now fix $u_2, \dots, u_\ell$ on the right hand side. Then to have equality, $\varphi_{e,3}$ cannot be a function of $\sigma_0 u_0$ or $\sigma_1 u_1$, so we have

$$\begin{aligned} \varphi_{e,3}(-, -, -, \sigma_2 u_2, \dots, \sigma_{\ell-1} u_{\ell-1}) = \\ \sigma_2 \varphi_{e,2}(-, -, u_2, \dots, u_\ell). \end{aligned}$$

Continuing this process in the same manner, we finally reduce the last equation, $j = \ell$, to

$$\begin{aligned} \varphi_{e,\ell}(-, -, \dots, -, \sigma_{\ell-1} u_{\ell-1}) = \\ \sigma_{\ell-1} \varphi_{e,\ell-1}(-, -, \dots, -, u_{\ell-1}, u_\ell). \end{aligned}$$

Summarizing our results, we can rewrite (33) as

$$\begin{aligned} \varphi_{e,j}(-, -, \dots, -, \sigma_{j-1} u_{j-1}, \dots, \sigma_{\ell-1} u_{\ell-1}) = \\ \sigma_{j-1} \varphi_{e,j-1}(-, -, \dots, -, u_{j-1}, \dots, u_\ell) \end{aligned} \quad (35)$$

for $j = 2, \dots, \ell$.

Equations (32) and (35) show that $\varphi_{e,j}$ is only a function of the last $\ell - j + 1$ components, for $j = 0, \dots, \ell$. We can formalize this information by using a new notation.

For $j = 0, \dots, \ell - 1$, define a function $\phi_j : \mathbf{U}_j \times \dots \times \mathbf{U}_\ell \rightarrow \mathbf{U}_j$ by

$$\phi_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \stackrel{\text{def}}{=} \varphi_{e,j}(-, -, \dots, -, \mathbf{u}_j, \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell),$$

and for $j = \ell$, define

$$\phi_\ell(\mathbf{u}_\ell) \stackrel{\text{def}}{=} \varphi_{e,\ell}(-, -, \dots, -, \mathbf{u}_\ell).$$

Then $\varphi_e = (\phi_0, \phi_1, \dots, \phi_\ell)$, and we can rewrite (32) and (35) as

$$\begin{aligned} \phi_j(\sigma_{j-1} \mathbf{u}_{j-1} | \sigma_j \mathbf{u}_j, \dots, \sigma_{\ell-1} \mathbf{u}_{\ell-1}) = \\ \sigma_{j-1} \phi_{j-1}(\mathbf{u}_{j-1} | \mathbf{u}_j, \dots, \mathbf{u}_\ell) \end{aligned} \quad (36)$$

for $j = 1, \dots, \ell - 1$, and

$$\phi_\ell(\sigma_{\ell-1} \mathbf{u}_{\ell-1}) = \sigma_{\ell-1} \phi_{\ell-1}(\mathbf{u}_{\ell-1} | \mathbf{u}_\ell) \quad (37)$$

for $j = \ell$. Note that (36)-(37) can be explained using commutative diagrams.

For $j = 0, \dots, \ell - 1$, for each fixed $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, the function $\phi_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ is a function $\phi_j : \mathbf{U}_j \times (\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \rightarrow \mathbf{U}_j$, essentially a function indexed by $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ with domain $\mathbf{U}_j$ and range a subset of $\mathbf{U}_j$. Note that when the range is all of $\mathbf{U}_j$, it is well defined to say $\phi_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ is 1-1 and onto. The following technical lemma is crucial, but the proof is straightforward using finite induction.

Lemma 36 $\varphi_e = (\phi_0, \dots, \phi_\ell)$ is 1-1 and onto if and only if $\phi_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ is 1-1 and onto for each fixed $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, for $j = 0, \dots, \ell - 1$, and ϕ_ℓ is 1-1 and onto.

Using Lemma 36, we are able to characterize an automorphism of $\mathcal{D}_\mathbf{U}$ as follows.

Theorem 37 $\varphi = (\varphi_s, \varphi_e)$ is an automorphism of $\mathcal{D}_\mathbf{U}$ if and only if $\varphi_e = (\phi_0, \phi_1, \dots, \phi_\ell)$ and $\varphi_s = (\phi_1, \dots, \phi_\ell)$, where $\phi_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ is 1-1 and onto for each fixed $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, for $j = 0, \dots, \ell - 1$, and ϕ_ℓ is 1-1 and onto, and (36)-(37) is satisfied for $j = 1, \dots, \ell$.

We can solve the set of equations (36)-(37) by starting with $j = \ell$ and working backwards. For $j = \ell - 1, \dots, 0$, for each $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, define the function $\beta_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ to be a permutation on $\mathbf{U}_j$ indexed by $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$, i.e., $\beta_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ is a 1-1 and onto function $\beta_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : \mathbf{U}_j \rightarrow \mathbf{U}_j$. We will construct permutation $\beta_j(\cdot | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : \mathbf{U}_j \rightarrow \mathbf{U}_j$ using permutations $\beta_{j,k}(\cdot | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow \mathbf{U}_j$, for $j \leq k \leq \ell$. If β_j is to be 1-1 and onto, then the images of $\beta_{j,k}$, for $j \leq k \leq \ell$, must be disjoint, and the union of the images must be all of $\mathbf{U}_j$. We will show that $\beta_{j,k}$ must have image $U_{j,k}$, or $\beta_{j,k}(\cdot | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow U_{j,k}$.

First, for $j = \ell$, pick any permutation β_ℓ on $\mathbf{U}_\ell$.

Next for $j = \ell - 1$, we must solve the equation

$$\sigma_{\ell-1} \beta_{\ell-1}(\mathbf{u}_{\ell-1} | \mathbf{u}_\ell) = \beta_\ell(\sigma_{\ell-1} \mathbf{u}_{\ell-1}), \quad (38)$$

where

$$\mathbf{u}_{\ell-1} \in (U_{\ell-1,\ell} U_{\ell-1,\ell-1})^T,$$

and

$$\sigma_{\ell-1} \mathbf{u}_{\ell-1} = \mathbf{u}_\ell \in (U_{\ell,\ell})^T.$$

We know that β_ℓ is 1-1 and onto, and so $\beta_\ell(\cdot) : U_{\ell,\ell} \rightarrow U_{\ell,\ell}$. If $\beta_{\ell-1}$ is to be 1-1 and onto, studying the commutative diagram given by (38) shows that $\beta_{\ell-1}$ must take $U_{\ell-1,\ell}$ to $U_{\ell-1,\ell}$, or $\beta_{\ell-1,\ell}(\cdot | \mathbf{u}_\ell) : U_{\ell-1,\ell} \rightarrow U_{\ell-1,\ell}$. Then we must have

$$\zeta_{\ell-1,\ell} \beta_{\ell-1,\ell}(u_{\ell-1,\ell} | \mathbf{u}_\ell) = \beta_{\ell,\ell}(\zeta_{\ell-1,\ell} u_{\ell-1,\ell}),$$

which can be written as

$$\beta_{\ell-1,\ell}(u_{\ell-1,\ell} | \mathbf{u}_\ell) = \beta_{\ell,\ell}(u_{\ell-1,\ell}).$$

Since $\beta_{\ell-1}$ takes $U_{\ell-1,\ell}$ to $U_{\ell-1,\ell}$, then we know $\beta_{\ell-1}$ must take $U_{\ell-1,\ell-1}$ to $U_{\ell-1,\ell-1}$. Then the permutation $\beta_{\ell-1,\ell-1}$ must satisfy $\beta_{\ell-1,\ell-1}(\cdot | \mathbf{u}_\ell) : U_{\ell-1,\ell-1} \rightarrow U_{\ell-1,\ell-1}$, but is otherwise arbitrary.

In general, for arbitrary j , $0 \leq j < \ell$, we have that $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow U_{j,k}$ is a permutation satisfying

$$\begin{aligned} \beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) = \\ \beta_{j+1,k}(u_{j,k} | \sigma_{j+1} \mathbf{u}_{j+1}, \dots, \sigma_{\ell-1} \mathbf{u}_{\ell-1}). \end{aligned}$$

for $j < k \leq \ell$, and $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ is arbitrary for $j = k$. In this way we obtain the following theorem, written as an algorithm.

Algorithm 38 Any solution of the set of equations (36)-(37) which is an automorphism of $\mathcal{D}_\mathbf{U}$ can be found as follows.

1. Pick any permutation β_ℓ on $\mathbf{U}_\ell$.

2.

FOR $j = \ell - 1, \dots, 0$ (working backwards in order),

FOR k satisfying $j < k \leq \ell$,

for each $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, let $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow U_{j,k}$ be the permutation on $U_{j,k}$ satisfying

$$\begin{aligned} \beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) = \\ \beta_{j+1,k}(u_{j,k} | \sigma_{j+1} \mathbf{u}_{j+1}, \dots, \sigma_{\ell-1} \mathbf{u}_{\ell-1}). \end{aligned} \quad (39)$$

ENDFOR

FOR $k = j$,

for each $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, let $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow U_{j,k}$ be any permutation on $U_{j,k}$.

ENDFOR

For each $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, define a permutation $\beta_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : \mathbf{U}_j \rightarrow \mathbf{U}_j$ using permutations $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow U_{j,k}$ for k such that $j \leq k \leq \ell$.

ENDFOR

3. Set $\varphi_e = (\beta_0, \beta_1, \dots, \beta_\ell)$, $\varphi_s = (\beta_1, \dots, \beta_\ell)$, and $\varphi = (\varphi_s, \varphi_e)$. Then φ is an automorphism of $\mathcal{D}_\mathbf{U}$, and the set of φ obtained this way is the automorphism group of $\mathcal{D}_\mathbf{U}$, $\text{Aut}(\mathcal{D}_\mathbf{U})$.

We now show the automorphism group can be described using less memory in the index $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$ of $\beta_j(\cdot | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell)$. For $0 \leq j \leq \ell$ and $0 \leq k \leq \ell$, we define a column $\mathbf{u}_{j,>k}$ in the following way. If $j > k$, then $\mathbf{u}_{j,>k} \stackrel{\text{def}}{=} \mathbf{u}_j$. If $j \leq k$, then $\mathbf{u}_j$ is of the form

$$\mathbf{u}_j = (u_{j,\ell} \ \cdots \ u_{j,k+1} \ u_{j,k} \ \cdots \ u_{j,j})^T,$$

and we define $\mathbf{u}_{j,>k}$ as

$$\mathbf{u}_{j,>k} \stackrel{\text{def}}{=} (u_{j,\ell} \ \cdots \ u_{j,k+1} \ 0 \ \cdots \ 0)^T.$$

In other words, the elements $u_{j,i}$ for $j \leq i \leq k$ are set to 0. (If $k = \ell$, then all elements are set to 0.) For $\mathbf{u}_j, \mathbf{u}'_j \in \mathbf{U}_j$, define an equivalence relation $\mathbf{u}_j \equiv \mathbf{u}'_j$ if and only if $\mathbf{u}_{j,>k} = \mathbf{u}'_{j,>k}$. Then for each fixed j, k such that $0 \leq j \leq \ell$, $0 \leq k \leq \ell$, the equivalence relation $\equiv$ defines a family of equivalence classes which partition $\mathbf{U}_j$.

In the following algorithm, we are interested in permutations $\beta_j(\cdot | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{\ell,>k})$, $0 \leq j < \ell$, with indices $(\mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{\ell,>k})$ having a memory of at most κ , where $\kappa \leq \ell - k$; that is indices of the form

$$(\mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell, \quad (40)$$

where $(\mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_{j+\kappa+1} \times \dots \times \mathbf{U}_\ell$, and κ satisfies $j+1 \leq j+\kappa \leq \ell$, with $\kappa \leq \ell - k$.

Algorithm 39 Any solution of the set of equations (36)-(37) which is an automorphism of $\mathcal{D}_\mathbf{U}$ can be found as follows.

1. Pick any permutation β_ℓ on $\mathbf{U}_\ell$.

2.

FOR $j = \ell - 1, \dots, 0$ (working backwards in order),

FOR k satisfying $j < k \leq \ell$,

for each $(\mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, let $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) : U_{j,k} \rightarrow U_{j,k}$ be the permutation on $U_{j,k}$ satisfying

$$\begin{aligned} &\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) = \\ &\beta_{j+1,k}(u_{j,k} | \sigma_{j+1} \mathbf{u}_{j+1,>k}, \dots, \sigma_{j+\kappa-1} \mathbf{u}_{j+\kappa-1,>k}, \mathbf{0}, \dots, \mathbf{0}). \end{aligned} \quad (41)$$

ENDFOR

FOR $k = j$,

for each $(\mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_{j+1} \times \dots \times$

$\mathbf{U}_\ell$, let $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) : U_{j,k} \rightarrow U_{j,k}$ be any permutation on $U_{j,k}$.

ENDFOR

For each $(\mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) \in \mathbf{U}_{j+1} \times \dots \times \mathbf{U}_\ell$, define a permutation $\beta_j(\mathbf{u}_j | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : \mathbf{U}_j \rightarrow \mathbf{U}_j$ using permutations $\beta'_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) : U_{j,k} \rightarrow U_{j,k}$ for k such that $j \leq k \leq \ell$, where

$$\begin{aligned} &\beta'_{j,k}(u_{j,k} | \mathbf{u}_{j+1}, \dots, \mathbf{u}_\ell) = \\ &\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}). \end{aligned}$$

ENDFOR

3. Set $\varphi_e = (\beta_0, \beta_1, \dots, \beta_\ell)$, $\varphi_s = (\beta_1, \dots, \beta_\ell)$, and $\varphi = (\varphi_s, \varphi_e)$. Then φ is an automorphism of $\mathcal{D}_\mathbf{U}$, and the set of φ obtained this way is the automorphism group of $\mathcal{D}_\mathbf{U}$, $\text{Aut}(\mathcal{D}_\mathbf{U})$.

Corollary 40 Any automorphism φ in $\text{Aut}(\mathcal{D}_\mathbf{U})$ is uniquely specified by the collection of permutations

$$\beta_{0,k}(\cdot | \mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}),$$

for each k such that $0 \leq k \leq \ell$, and each $(\mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_1 \times \dots \times \mathbf{U}_\ell$, with $\kappa = \ell - k$.

We call the collection of permutations in Corollary 40 *separating permutations*. Algorithm 39 shows that permutations in the graph automorphism group can be separated by rows and by columns. Corollary 40 shows that only the permutations $\beta_{0,k}$, $0 \leq k \leq \ell$, are required to generate the automorphism group.

Using the separating permutations, we can count how many distinct automorphisms there are. The symmetric group on $|U_{0,k}|$ objects is denoted $\mathbb{S}_{|U_{0,k}|}$.

Theorem 41 $|\text{Aut}(\mathcal{D}_\mathbf{U})|$ is given by

$$|\text{Aut}(\mathcal{D}_\mathbf{U})| = \prod_{k=0}^{\ell} |\mathbb{S}_{|U_{0,k}|}|^{\Upsilon_k}, \quad (42)$$

where

$$\Upsilon_k = \prod_{k'=k+1}^{\ell} |U_{0,k'}|^{(k'-k)}$$

for $k < \ell$ and $\Upsilon_k = 1$ for $k = \ell$.

Proof. Each distinct automorphism φ uses a set of separating permutations

$$\begin{aligned} &F_k = \{\beta_{0,k}(\cdot | \mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) | \\ &\text{index } (\mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_1 \times \dots \times \mathbf{U}_\ell\} \end{aligned}$$

for each k , $0 \leq k \leq \ell$. There is one permutation $\beta_{0,k}$ in the set F_k for each index $(\mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) \in \mathbf{U}_1 \times \dots \times \mathbf{U}_\ell$. Let $< F_k >$ be the number of distinct sets F_k that can be formed. Then

$$|\text{Aut}(\mathcal{D}_\mathbf{U})| = \prod_{k=0}^{\ell} < F_k >.$$

We now count the number of distinct sets F_k . For each fixed index $(\mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0})$ in F_k , there are $|\mathbb{S}_{|U_{0,k}|}|$ possible choices for permutation $\beta_{0,k}$. The number of possible indices $(\mathbf{u}_{1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0})$ in F_k is

$$\Upsilon_k = \prod_{k'=k+1}^{\ell} |U_{0,k'}|^{(k'-k)}$$

for $k < \ell$. For $k = \ell$, the number of possible indices is 1, so let $\Upsilon_\ell = 1$. Then

$$|F_k| = |\mathbb{S}_{|U_{0,k}|}|^{\Upsilon_k},$$

for $0 \leq k \leq \ell$. •

7. BASIS AND BRANCH GROUP STRUCTURE

7.1 Basis and structure

We define a basis using the natural shift structure of B . The groups in $\{X_j\}$ are normal in B . Thus we can form the sequence of quotient groups

$$X_0, X_1/X_0, \dots, X_j/X_{j-1}, \dots, X_\ell/X_{\ell-1}. \quad (43)$$

This gives a coset decomposition chain of B . Pick a coset representative x_j from each coset of X_j/X_{j-1} , for $j = 0, \dots, \ell$. (Note: we mean x_j is an element of B , not a coset in X_j/X_{j-1} .) This forms a complete system of coset representatives, or *basis* $\mathcal{B}$. It is well known that any element $g \in B$ can be uniquely represented by a vector $\mathbf{x}$ of coset representatives in a complete system of coset representatives $\mathcal{B}$,

$$\mathbf{x} = (x_0, x_1, \dots, x_\ell), \quad (44)$$

called an *expansion* of g , where

$$g = \prod_{j=0}^{\ell} x_j. \quad (45)$$

Since there are many ways to pick the complete system of coset representatives to form a basis $\mathcal{B}$, the expansion of g in (44) and product terms in (45) are not unique. We refer to two different bases as a *change of basis*.

For each $g \in B$, let *expansion set* S_g be the set of expansions of g using a complete system of coset representatives for each possible basis $\mathcal{B}$. Thus if $\mathcal{B}$ is any basis and $\mathbf{x} = (x_0, \dots, x_\ell)$ is the expansion of g in terms of the coset representatives of that basis $\mathcal{B}$, so that $g = \prod_{j=0}^{\ell} x_j$, then $\mathbf{x} \in S_g$. In another basis $\mathcal{B}'$, the expansion of g might be $\mathbf{x}' = (x'_0, \dots, x'_\ell)$ so that $g = \prod_{j=0}^{\ell} x'_j$, and then $\mathbf{x}' \in S_g$. Thus S_g is an equivalence class.

Theorem 42 Vector $\mathbf{x} = (x_0, \dots, x_\ell) \in S_g$ if and only if $x_j \in X_j$ and $g = \prod_{j=0}^{\ell} x_j$.

Proof. First assume $x_j \in X_j$ and $g = \prod_{j=0}^{\ell} x_j$. If $x_j \in X_j$, then x_j is a coset representative in X_j/X_{j-1} . The coset representatives x_j , $j = 0, \dots, \ell$, define a unique $g \in B$, given by $g = \prod_{j=0}^{\ell} x_j$. Then there is some basis $\mathcal{B}$ containing coset representatives x_j , $j = 0, \dots, \ell$, in a complete set of coset representatives, and $\mathbf{x} \in S_g$.

Now assume $\mathbf{x} \in S_g$. If $\mathbf{x} \in S_g$ then $g = \prod_{j=0}^{\ell} x_j$ and x_j are coset representatives in X_j/X_{j-1} , $j = 0, \dots, \ell$. Then $x_j \in X_j$. •

We will now show the set of expansion classes S_g , for each $g \in B$, forms a group. Consider any $\mathbf{x} \in S_g$ and any $\bar{\mathbf{x}} \in S_{\bar{g}}$. We define a product operation $\mathbf{x} \otimes \bar{\mathbf{x}}$ as

$$\begin{aligned} \mathbf{x} \otimes \bar{\mathbf{x}} &= (x_0, \dots, x_\ell) \otimes (\bar{x}_0, \dots, \bar{x}_\ell) \\ &\stackrel{\text{def}}{=} (x_0 x_0^*, \dots, x_j x_j^*, \dots, x_\ell x_\ell^*), \end{aligned} \quad (46)$$

where x_j^* is defined by

$$x_j^*(x_{j+1} \cdots x_\ell) = (x_{j+1} \cdots x_\ell) \bar{x}_j, \quad (47)$$

for $0 \leq j < \ell$, and $x_\ell^* = \bar{x}_\ell$. The group operation is similar to a direct product operation but with a “twist”.

Let $\mathbf{x}' \stackrel{\text{def}}{=} (x_0 x_0^*, \dots, x_j x_j^*, \dots, x_\ell x_\ell^*)$. Fix j such that $0 \leq j < \ell$. Since X_j is normal in $X_{j+1}, \dots, X_\ell$, we know that $X_j(x_{j+1} \cdots x_\ell) = (x_{j+1} \cdots x_\ell) X_j$. And since $\bar{x}_j \in X_j$, then from (47) we know $x_j^* \in X_j$. Then $x_j x_j^* \in X_j$ and from Theorem 42, $\mathbf{x}' \in S_{g'}$ where $g' = \prod_{j=0}^{\ell} x_j x_j^*$.

We now show $g' = g\bar{g}$. We have

$$\begin{aligned} g' &= \prod_{j=0}^{\ell} x_j x_j^* \\ &= \left(\prod_{j=0}^{\ell} x_j \right) \left(\prod_{j=0}^{\ell} \bar{x}_j \right) \\ &= g\bar{g}. \end{aligned}$$

Thus for any $\mathbf{x} \in S_g$ and any $\bar{\mathbf{x}} \in S_{\bar{g}}$, we have shown $\mathbf{x} \otimes \bar{\mathbf{x}} \in S_{g\bar{g}}$.

We can improve on this result as follows.

Lemma 43 Fix any $\mathbf{x} \in S_g$. Define $\mathbf{x} \otimes S_{\bar{g}}$ to be the set of products $\{\mathbf{x} \otimes \bar{\mathbf{x}} | \bar{\mathbf{x}} \in S_{\bar{g}}\}$. Then $\mathbf{x} \otimes S_{\bar{g}} = S_{g'}$ and there is a 1-1 and onto map $\eta : \mathbf{x} \otimes S_{\bar{g}} \rightarrow S_{g'}$.

Proof. We have just shown that for any $\bar{\mathbf{x}} \in S_{\bar{g}}$, $\mathbf{x} \otimes \bar{\mathbf{x}} \in S_{g'}$. Thus we have $\mathbf{x} \otimes S_{\bar{g}} \subset S_{g'}$. It remains to show the map η is 1-1 and onto. But if $(\bar{x}_0, \dots, \bar{x}_\ell) \neq (\tilde{x}_0, \dots, \tilde{x}_\ell)$, it is clear from (47) that $\mathbf{x} \otimes \bar{\mathbf{x}} \neq \mathbf{x} \otimes \tilde{\mathbf{x}}$. Thus the map η is 1-1. We now show η is onto. Pick any $\mathbf{x}' = (x'_0, \dots, x'_\ell) \in S_{g'}$. Find $(\hat{x}_0, \dots, \hat{x}_\ell)$ such that

$$(x'_0, \dots, x'_\ell) = (x_0 \hat{x}_0, \dots, x_\ell \hat{x}_\ell). \quad (48)$$

We can rewrite this as

$$\begin{aligned} (x'_0, \dots, x'_\ell) &= (x_0 \hat{x}_0, \dots, x_\ell \hat{x}_\ell) \\ &= (x_0, \dots, x_\ell) \otimes (\hat{x}_0, \dots, \hat{x}_\ell), \end{aligned}$$

where $\hat{x}_j$ is defined by

$$\dot{x}_j(x_{j+1} \cdots x_\ell) = (x_{j+1} \cdots x_\ell)\hat{x}_j, \quad (49)$$

for $j = 0, \dots, \ell - 1$, and $\hat{x}_\ell = \dot{x}_\ell$. Then we have

$$\begin{aligned} g' &= \prod_{j=0}^{\ell} x'_j \\ &= \prod_{j=0}^{\ell} x_j \dot{x}_j \\ &= \left(\prod_{j=0}^{\ell} x_j \right) \left(\prod_{j=0}^{\ell} \hat{x}_j \right). \end{aligned}$$

But since $g' = g\bar{g}$ and $\prod_{j=0}^{\ell} x_j = g$, we must have $\prod_{j=0}^{\ell} \hat{x}_j = \bar{g}$. It is clear from (48) that $\dot{x}_j \in X_j$, and then from (49) that $\hat{x}_j \in X_j$. Thus $\hat{\mathbf{x}} \stackrel{\text{def}}{=} (\hat{x}_0, \dots, \hat{x}_\ell) \in S_{\bar{g}}$, and there exists $\hat{\mathbf{x}} \in S_{\bar{g}}$ such that $\mathbf{x} \otimes \hat{\mathbf{x}} = \mathbf{x}'$. This means map η is onto. •

Lemma 44 Fix any $\bar{\mathbf{x}} \in S_{\bar{g}}$. Define $S_g \otimes \bar{\mathbf{x}}$ to be the set of products $\{\mathbf{x} \otimes \bar{\mathbf{x}} | \mathbf{x} \in S_g\}$. Then $S_g \otimes \bar{\mathbf{x}} = S_{g'}$ and there is a 1-1 and onto map $\eta : S_g \otimes \bar{\mathbf{x}} \rightarrow S_{g'}$.

Proof. Similar to above proof. •

Let S be the set of expansion classes, $S = \cup_{g \in B} S_g$. In view of Lemmas 43 and 44, we have the following.

Theorem 45 For $g, \bar{g} \in B$, define the product operation $S_g \otimes S_{\bar{g}}$ on sets in S to be the set of products $\{\mathbf{x} \otimes \bar{\mathbf{x}} | \mathbf{x} \in S_g, \bar{\mathbf{x}} \in S_{\bar{g}}\}$. In view of Lemmas 43 and 44, the product operation $S_g \otimes S_{\bar{g}}$ is well defined and

$$S_{g'} = \{\mathbf{x} \otimes \bar{\mathbf{x}} | \mathbf{x} \in S_g, \bar{\mathbf{x}} \in S_{\bar{g}}\},$$

where $g' = g\bar{g}$.

Consider $S_h \otimes (S_g \otimes S_{\bar{g}})$ and $(S_h \otimes S_g) \otimes S_{\bar{g}}$. We have

$$\begin{aligned} S_h \otimes (S_g \otimes S_{\bar{g}}) &= S_h \otimes S_{g\bar{g}} \\ &= S_{h(g\bar{g})} \\ &= S_{(hg)\bar{g}} \\ &= S_{hg} \otimes S_{\bar{g}} \\ &= (S_h \otimes S_g) \otimes S_{\bar{g}}. \end{aligned}$$

Thus the product operation “ $\otimes$ ” on S is associative because B is associative.

Then we can form an *expansion class group* S_B on elements of S using group operation $S_g \otimes S_{\bar{g}}$ for $S_g, S_{\bar{g}} \in S$. The identity element of S_B is S_1 , and the inverse of S_g is $S_{g^{-1}}$.

Theorem 46 The set S of expansion classes S_g , for each $g \in B$, along with product operation $\otimes$, forms an expansion class group S_B .

Theorem 47 The bijection $\lambda : B \rightarrow S_B$ given by the assignment $\lambda : g \mapsto S_g$ is an isomorphism from branch group B to class group S_B , that is

$$S_{g\bar{g}} = S_g \otimes S_{\bar{g}}.$$

Note that group multiplication in S_B , given by (46)-(47), is local, causal, and iterative.

If B is abelian, it can be seen from (47) that $x_j^* = \bar{x}_j$ for $j = 0, \dots, \ell$. Then the product operation $\mathbf{x} \otimes \bar{\mathbf{x}}$ in (46) becomes

$$\begin{aligned} \mathbf{x} \otimes \bar{\mathbf{x}} &= (x_0, \dots, x_\ell) \otimes (\bar{x}_0, \dots, \bar{x}_\ell) \\ &= (x_0 \bar{x}_0, \dots, x_j \bar{x}_j, \dots, x_\ell \bar{x}_\ell). \end{aligned}$$

Thus in the abelian case there is no “twist” in the group operation “ $\otimes$ ”, and class group S_B has properties of a direct product.

7.2 Change of generator basis

In the previous subsection, we introduced a vector basis, in which any element $g \in B$ can be uniquely represented by a vector $\mathbf{x}$ of coset representatives of (43),

$$\mathbf{x} = (x_0, x_1, \dots, x_\ell).$$

The controllable matrix gives a collection of coset representatives $\{r_{j,k}\}$ for $0 \leq k \leq \ell$ and $0 \leq j \leq k$. The product $x_j = \prod_{k=j}^{\ell} r_{j,k}$ is a representative of a coset in X_j/X_{j-1} . Therefore we can define a vector $\mathbf{x}$ of coset representatives of (43) using

$$\mathbf{x} = \left(\left(\prod_{k=0}^{\ell} r_{0,k} \right), \left(\prod_{k=1}^{\ell} r_{1,k} \right), \dots, \left(\prod_{k=j}^{\ell} r_{j,k} \right), \dots, (r_{\ell,\ell}) \right).$$

We call the basis formed this way a *generator basis*. In general there are vector bases which cannot be formed using a generator basis; therefore the generator bases form a strict subset of the vector bases. However the generator bases can be used to form a minimal encoder [3], as given in [3] and Section 5. In this subsection and the next we only consider generator bases.

Consider using two generator bases to form two encoders. Since generator bases give minimal encoders, the graphs of the two encoders are isomorphic [3]. In this section we study how the graph automorphism group places restrictions on the change of generator basis.

We have just seen that an edge e of the shift register graph $\mathcal{D}_{\mathbf{U}}$ can be uniquely identified by a set $\{u_{j,k}\}$ of integers, where each $u_{j,k} \in U_{j,k}$, $0 \leq j \leq k$, $0 \leq k \leq \ell$. Clearly there is a 1-1 correspondence between $U_{j,k} \in \mathbf{U}$ in (18) and $Q_{j,k}$ in (16), such that $u_{j,k} \in U_{j,k}$ corresponds to a single coset $q_{j,k}(u_{j,k})$ in quotient group $Q_{j,k}$, for $0 \leq j \leq k$, $0 \leq k \leq \ell$. (The argument $u_{j,k}$ in $q_{j,k}(u_{j,k})$ is regarded as an index for the cosets $q_{j,k}$ in $Q_{j,k}$.) When we select a generator basis $\mathcal{B}$, we select a representative $r_{j,k}(u_{j,k}) \in q_{j,k}(u_{j,k})$ for each coset $q_{j,k}(u_{j,k})$ in $Q_{j,k}$. The product of the representatives gives a unique $g \in B$, which is

the label of edge e in the encoder $\mathcal{D}_{\mathbf{U},E}$ determined by generator basis $\mathcal{B}$. Then we have the correspondences:

$$e \longleftrightarrow \{u_{j,k}\} \longleftrightarrow \{q_{j,k}(u_{j,k})\} \longleftrightarrow \{r_{j,k}(u_{j,k})\} \longleftrightarrow g = \prod r_{j,k}(u_{j,k}). \quad (50)$$

Suppose we have another generator basis $\mathcal{B}'$. Then we select a representative $r'_{j,k}(u'_{j,k}) \in q_{j,k}(u_{j,k})$ for each coset $q_{j,k}(u_{j,k})$ in $Q_{j,k}$, where we may not have $r_{j,k}(u_{j,k}) = r'_{j,k}(u_{j,k})$. The basis $\mathcal{B}'$ gives an encoder $\mathcal{D}_{\mathbf{U},E'}$, and for an edge e' in $\mathcal{D}_{\mathbf{U},E'}$, we have the correspondences:

$$e' \longleftrightarrow \{u'_{j,k}\} \longleftrightarrow \{q_{j,k}(u'_{j,k})\} \longleftrightarrow \{r'_{j,k}(u'_{j,k})\} \longleftrightarrow g' = \prod r'_{j,k}(u'_{j,k}). \quad (51)$$

The two generator bases $\mathcal{B}$ and $\mathcal{B}'$ give encoders $\mathcal{D}_{\mathbf{U},E}$ and $\mathcal{D}_{\mathbf{U},E'}$, respectively. Since the encoders are minimal, they must be graph isomorphic [3]. This means there is a graph isomorphism φ which takes edge e with label g in $\mathcal{D}_{\mathbf{U},E}$ to edge e' with label g' in $\mathcal{D}_{\mathbf{U},E'}$, where $g = g'$. Then we have the relations:

$$\begin{array}{ccccccc} e & \longleftrightarrow & \{u_{j,k}\} & \longleftrightarrow & \{q_{j,k}(u_{j,k})\} & \longleftrightarrow & \{r_{j,k}(u_{j,k})\} \longleftrightarrow g = \prod r_{j,k}(u_{j,k}) \\ \downarrow \varphi & & & & & & \downarrow \varphi \\ e' & \longleftrightarrow & \{u'_{j,k}\} & \longleftrightarrow & \{q_{j,k}(u'_{j,k})\} & \longleftrightarrow & \{r'_{j,k}(u'_{j,k})\} \longleftrightarrow g' = \prod r'_{j,k}(u'_{j,k}) = g. \end{array} \quad (52)$$

Again, note that there is a representative $r_{j,k}(u'_{j,k})$ in $\mathcal{B}$ of coset $q_{j,k}(u'_{j,k})$, which may not be the same as representative $r'_{j,k}(u'_{j,k})$ in $\mathcal{B}'$ of coset $q_{j,k}(u'_{j,k})$.

But we know the isomorphism φ gives a bijection $\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) : U_{j,k} \rightarrow U_{j,k}$. Then we must have

$$\beta_{j,k}(\cdot | \mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0}) : u_{j,k} \mapsto u'_{j,k}.$$

But we know that g uniquely identifies the cosets $q_{j,k}(u_{j,k})$, and therefore the integers $u_{j,k} \in U_{j,k}$. Thus g specifies the index $(\mathbf{u}_{j+1,>k}, \dots, \mathbf{u}_{j+\kappa,>k}, \mathbf{0}, \dots, \mathbf{0})$, and we show this dependency by writing $(\mathbf{u}_{j+1,>k}^g, \dots, \mathbf{u}_{j+\kappa,>k}^g, \mathbf{0}, \dots, \mathbf{0})$. This means

$$g = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r_{j,k}(u_{j,k}) \quad (53)$$

$$= \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r'_{j,k}(\beta_{j,k}(u_{j,k} | \mathbf{u}_{j+1,>k}^g, \dots, \mathbf{u}_{j+\kappa,>k}^g, \mathbf{0}, \dots, \mathbf{0})) \quad (54)$$

$$= \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r'_{j,k}(u'_{j,k}). \quad (55)$$

Thus we can represent g in basis $\mathcal{B}$ using (53), and in basis $\mathcal{B}'$ using (55). The assignment $r_{j,k}(u_{j,k}) \mapsto r'_{j,k}(u'_{j,k})$ gives the change of basis. This proves a change of basis theorem.

Theorem 48 *Let $\mathcal{B}$ and $\mathcal{B}'$ be two generator bases with representatives $\{r_{j,k}\}$ and $\{r'_{j,k}\}$, respectively. Let $g \in B$ be represented in generator basis $\mathcal{B}$ as*

$$g = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r_{j,k}(u_{j,k}).$$

Then the representation of g in generator basis $\mathcal{B}'$ is

$$g = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r'_{j,k}(u'_{j,k}),$$

where the map

$$r_{j,k}(u_{j,k}) \mapsto r'_{j,k}(u'_{j,k}) \quad (56)$$

is found using

$$\beta_{j,k}(\cdot | \mathbf{u}_{j+1,>k}^g, \dots, \mathbf{u}_{j+\kappa,>k}^g, \mathbf{0}, \dots, \mathbf{0}) : u_{j,k} \mapsto u'_{j,k}, \quad (57)$$

and $\beta_{j,k}$ is from the graph isomorphism φ taking label g with edge representation e in $\mathcal{B}$ to label g with edge representation e' in $\mathcal{B}'$.

We can think of the change of basis theorem as relating two coset decomposition chains. For a particular g , there is a bijective mapping between representatives in one coset chain to representatives in the other depending on a memory given by the index $(\mathbf{u}_{j+1,>k}^g, \dots, \mathbf{u}_{j+\kappa,>k}^g, \mathbf{0}, \dots, \mathbf{0})$, for $0 \leq j \leq k$, $0 \leq k \leq \ell$, as shown in (56)-(57). For any other element $h \in B$, the mapping from representatives in

one coset chain to the other is the same as for g for those j, k such that $(\mathbf{u}_{j+1, > k}^g, \dots, \mathbf{u}_{j+\kappa, > k}^g, \mathbf{0}, \dots, \mathbf{0}) = (\mathbf{u}_{j+1, > k}^h, \dots, \mathbf{u}_{j+\kappa, > k}^h, \mathbf{0}, \dots, \mathbf{0})$. If the graph automorphism for the change of basis is independent of index $(\mathbf{u}_{j+1, > k}^g, \dots, \mathbf{u}_{j+\kappa, > k}^g, \mathbf{0}, \dots, \mathbf{0})$ for all g and all $0 \leq j \leq k$, $0 \leq k \leq \ell$, there is a bijective mapping between representatives in one coset chain to the other which is independent of g . Thus there is a fixed 1-1 mapping of the two bases $\mathcal{B}$ and $\mathcal{B}'$ which gives the representation of g in $\mathcal{B}'$ from the representation of g in $\mathcal{B}$, independent of g . Any de Bruijn shift register is an example of this, such as a binary convolutional code.

We have just seen that the change of generator basis, the branch group, the shift register graph, and the graph automorphism group are intimately related. This is shown by the following theorem.

Theorem 49 *The number of generator bases is upper bounded by $|\text{Aut}(\mathcal{D}_{\mathbf{U}})|$, where $|\text{Aut}(\mathcal{D}_{\mathbf{U}})|$ is given in (42).*

Proof. A generator basis gives a minimal encoder. The graphs of any two minimal encoders are related by a graph isomorphism [3]. •

7.3 Symmetry and duality for abelian B

In this paragraph, we give a brief review of the minimal encoder construction of Forney and Trott (see *Encoder Construction*, p. 1509 of [3]). **In this paragraph, the notation in [3] is used.** The j -controllable subcode C_j of a group code C is defined as the set of combinations of code sequences of span $j+1$ or less:

$$C_j = \prod_{k \in \mathbb{Z}} C_{[k, k+j]}.$$

They show

$$C_0 \subseteq C_1 \subseteq \dots C_{j-1} \subseteq C_j \subseteq \dots C_\nu = C$$

is a normal series. Then in their *Code Granule Theorem*, they show C_j/C_{j-1} is isomorphic to the direct product $\prod_{k \in \mathbb{Z}} \Gamma_{[k, k+j]}$, where $\Gamma_{[k, k+j]}$ is a granule, a particular quotient group. Then it follows (p. 1509) that the set $\prod_{k \in \mathbb{Z}} [\Gamma_{[k, k+j]}]$ is a set of coset representatives for the cosets of C_{j-1} in C_j , where $[\Gamma_{[k, k+j]}] = \{\mathbf{c}(\gamma_{k,j})\}$ is a set of coset representatives for the granule $\Gamma_{[k, k+j]}$. This means (*Generator Theorem*) that every code sequence $\mathbf{c}$ can be uniquely expressed as a product

$$\mathbf{c} = \prod_{0 \leq j \leq \nu} \prod_{k \in \mathbb{Z}} \mathbf{c}(\gamma_{k,j}) \quad (58)$$

of generators $\mathbf{c}(\gamma_{k,j})$.

We have seen in Section 4 that the generators used by Forney and Trott are the same as the generators used in the encoder here. We can then rewrite (58) using our notation. The time index k in [3] corresponds to time

index t in our notation. The time index j in [3] corresponds to k in our notation. Thus the generator $\mathbf{c}(\gamma_{k,j})$ in (58) corresponds to $\mathbf{g}_{t,k}$ in our notation. Further ν in [3] corresponds to ℓ here. Then we can rewrite (58) in our notation as

$$\mathbf{c} = \prod_{0 \leq k \leq \ell} \prod_{t \in \mathbb{Z}} \mathbf{g}_{t,k}. \quad (59)$$

The output at time t , or branch b'_t in the branch group version of the encoder, is given by

$$b'_t = \prod_{k=0}^{\ell} \left(\prod_{j=k}^0 \chi_t(\mathbf{g}_{t-j,k}) \right).$$

For our purposes, it is more convenient to consider the time reversed version of (59). This is also a valid encoder since the argument in [3] is symmetric with respect to time reversal. Then the output at time t for the time reversed encoder, branch b''_t , is given by

$$b''_t = \prod_{k=0}^{\ell} \left(\prod_{j=0}^k \chi_t(\mathbf{g}_{t-j,k}) \right). \quad (60)$$

Using (20) we can rewrite (60) as

$$b''_t = \prod_{k=0}^{\ell} \left(\prod_{j=0}^k r_{j,k} \right). \quad (61)$$

We now compare (61) and the encoder (19) in Section 5. In (19), for fixed j the inner product (the term in parentheses of (19)) is a product of terms in a single column of the controllable matrix, and the outer product can be considered to be a row product. Then we refer to encoder (19) as a *column-row* encoder. In the Forney and Trott encoder (61), for fixed k the inner product (the term in parentheses of (61)) is a product of terms in a single row of the controllable matrix, and the outer product can be considered to be a column product. Thus we refer to the Forney and Trott encoder as a *row-column* encoder. This terminology points out a distinct difference between the two encoders. However note that one encoder can be transformed to the other by merely interchanging the inner and outer product.

Fix a generator basis $\mathcal{B}^*$. This gives a set of representatives $\{r_{j,k}^*\}$. Using basis $\mathcal{B}^*$, and emphasizing the connection to an underlying graph $\mathcal{D}_{\mathbf{U}}$, we rewrite (19) as

$$g = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r_{j,k}^*(u_{j,k}). \quad (62)$$

We know that we can use the same generator basis $\mathcal{B}^*$ for the Forney-Trott encoder. Then we have

$$g' = \prod_{k=0}^{\ell} \prod_{j=0}^k r_{j,k}^*(u_{j,k}). \quad (63)$$

In general we do not have $g = g'$ if the product terms in (62) and (63) are the same. But since both encoders are minimal, we know there is a graph automorphism φ containing $\beta_{j,k}$, where

$$\beta_{j,k}(\cdot | \mathbf{u}_{j+1, > k}, \dots, \mathbf{u}_{j+\kappa, > k}, \mathbf{0}, \dots, \mathbf{0}) : u_{j,k} \mapsto u'_{j,k},$$

such that

$$g = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r_{j,k}^*(u_{j,k}) = \prod_{j=0}^{\ell} \prod_{k=j}^{\ell} r_{j,k}^*(u'_{j,k}), \quad (64)$$

as in (53)-(55). In general (64) gives commutative restrictions on the basis representatives since the same representatives appear in different order in both encoders. And these restrictions can be obtained for each choice of basis $\mathcal{B}^*$. Therefore in general a nonabelian branch group B must satisfy extensive commutative restrictions on its elements.

Now assume B is abelian and both encoders use the same basis $\mathcal{B}$. Since B is abelian the product terms in (62) can be rearranged to form (63) so that $g = g'$. Then the graph isomorphism φ is trivial, and the states of the Forney-Trott encoder are the same as the states of the encoder here. This gives the following result.

Theorem 50 *Assume B is abelian. Fix any generator basis $\mathcal{B}$. Let $\mathcal{B}$ have representatives $\{r_{j,k}\}$. Then for any $g \in B$, we have*

$$g = \prod_{j=0}^{\ell} \left(\prod_{k=j}^{\ell} r_{j,k} \right) = \prod_{k=0}^{\ell} \left(\prod_{j=0}^k r_{j,k} \right). \quad (65)$$

The middle term describes the encoder given here, and the right term describes the branch group version of the Forney-Trott encoder. The middle term can be regarded as a time domain representation of g , and the right term as a spectral domain representation of g . Theorem 50 shows the two domain representations are equal, and there is a symmetry and duality between the time domain and spectral domain. We can rewrite the middle term of (65) as

$$\left(\prod_{k=0}^{\ell} r_{0,k} \right) \left(\prod_{k=1}^{\ell} r_{1,k} \right) \cdots \left(\prod_{k=j}^{\ell} r_{j,k} \right) \cdots (r_{\ell,\ell}). \quad (66)$$

Then Theorem 50 shows that the time domain representation can be transformed to the spectral domain by using the row product in (66), similar to a trace. Note that each parentheses term in (66) is a component of the vector basis.

References

[1] B. Kitchens, “Expansive dynamics on zero-dimensional groups,” *Ergodic Theory and Dynamical Systems* **7**, pp. 249-261, 1987.

[2] J. C. Willems, “Models for dynamics,” in *Dynamics Reported*, vol. 2, U. Kirchgraber and H. O. Walther, Eds., New York: John Wiley, 1989.

[3] G. D. Forney, Jr. and M. D. Trott, “The dynamics of group codes: state spaces, trellis diagrams, and canonical encoders,” *IEEE Trans. Inform. Theory*, vol. 39, pp. 1491-1513, Sept. 1993.

[4] H.-A. Loeliger and T. Mittelholzer, “Convolutional codes over groups,” *IEEE Trans. Inform. Theory, Part I*, vol. 42, pp. 1660-1686, Nov. 1996.

[5] M. D. Trott, “The algebraic structure of trellis codes,” Ph.D. thesis, Stanford Univ., Aug. 1992.

[6] G. D. Forney, Jr., “Geometrically uniform codes,” *IEEE Trans. Inform. Theory*, vol. 37, pp. 1241-1260, Sept. 1991.

[7] G. D. Forney, Jr., “Minimal realizations of linear systems: the “shortest basis” approach,” arXiv:0910.4336v3 [cs.IT], 24 Aug 2010.

[8] J. P. Sarvis and M. D. Trott, “Useful groups for trellis codes,” in *Proc. IEEE Int. Symp. Inform. Theory*, Whistler, BC, Canada, Sept. 17-22, 1995, p. 308.

[9] N. T. Sindhusayana, B. Marcus, and M. Trott, “Homogeneous shifts,” *IMA J. Math. Contr. Inform.*, vol. 14, pp. 255-287, 1997.

[10] J. J. Rotman, *An Introduction to the Theory of Groups* (4th edition), Springer, New York, 1995.

[11] M. Hall, Jr., *The Theory of Groups*, Chelsea, New York, 1959.

[12] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge Univ. Press, New York, 1995.

[13] J. Dénes and A. D. Keedwell, *Latin squares and Their Applications*, Budapest, Akadémiai Kiado, 1974.

[14] D. Jungnickel, “Latin squares, their geometries and their groups. A survey,” in *Coding Theory and Design Theory, Part II* (D. Ray-Chaudhuri, ed.), vol. 21 of *IMA Volumes in Mathematics and its Applications*, pp. 166-225, Springer, 1992.

[15] K. M. Mackenthun, Jr., “On groups with a shift structure: the Schreier matrix and an algorithm,” in *41st Annual Conf. on Information Sciences and Systems*, Baltimore, MD, March 14-16, 2007.

[16] K. M. Mackenthun, Jr., “Group codes and the Schreier matrix form,” arXiv:1106.3981v1 [cs.IT], 20 Jun 2011.